

Of this form, of which an immature female is before me, one can say nothing except that, for all evidence to the contrary, it may well be the young of some species of *Eurypelma* or other closely allied genus. It is especially to be borne in mind that the circular form of the sternum is characteristic of immature specimens of most Eurypelmiform spiders.

### CRATORRHAGUS.

*Cratorrhagus*, E. Simon, Act. Soc. Linn. Bord. xliv. p. 330 (1892).

Type *Cratorrhagus concolor*, Simon.

#### ***Cratorrhagus cervinus*.** (Tab. II. figg. 1, 1 *a-c*, ♀.)

*Cratorrhagus cervinus*, E. Simon, Act. Soc. Linn. Bord. xliv. p. 330 (1892)<sup>1</sup>.

Type, ♀, in coll. E. Simon, Paris.

*Hab.* MEXICO<sup>1</sup>.

The type of *C. cervinus*, which is now before me, is certainly not congeneric with a specimen of *C. concolor* from the Mediterranean, so far as one can judge from an immature specimen. But I would not hazard an opinion with specimens not yet adult, and I hesitate therefore to tabulate the characters of the female from Mexico.

### MYGALARACHNE.

*Mygalarachne*, Ausserer, Verh. zool.-bot. Ges. Wien, xxi. p. 206 (1871).

#### ***Mygalarachne brevipes*.**

*Mygalarachne brevipes*, Auss. loc. cit. p. 206<sup>1</sup>; Simon, Act. Soc. Linn. Bord. xliv. p. 339<sup>2</sup>.

Type, ♀, in Vienna Museum.

*Hab.* HONDURAS<sup>1 2</sup>.

Without opportunity of examining the type I am unable to offer any opinion as to the systematic position of this spider. Neither does Ausserer give any descriptive clue as to its identity, save only "die Sammtbürste am innerrande der Schenkel fehlt."

### DAVUS.

*Davus*, O. P.-Cambridge, Biol. Centr.-Amer., Arachn. Aran. i. p. 91 (1892).

Femora iv. not scopulate on the inner side. Protarsi i. and ii. scopulate beneath in their apical half only.

Protarsus iv. not scopulate, very spinose. Tibia i. with two short spurs at the apex beneath.

#### ***Davus fasciatus*.** (Tab. III. figg. 4, 4 *a-d*, ♂.)

*Davus fasciatus*, O. P.-Cambr. Biol. Centr.-Amer., Arachn. Aran. i. p. 91 (♂)<sup>1</sup>.

*Pella fasciata*, O. P.-Cambr. loc. cit. t. 12. figg. 4, 4 *a-d* (♂).

*Hab.* COSTA RICA (*Boucard*<sup>1</sup>).

The type of this genus and species is a male, dry, and rather the worse for wear. I  
BIOL. CENTR.-AMER., Arachn. Aran., Vol. II., April 1899. 6‡