

## B. Lower margin of fang-groove with 4 teeth.

a. Eyes more closely grouped. Anterior eyes equidistant. Anterior row strongly procurved. No small central apical protarsal spine . . . . . *bulbosa*.

b. Eyes widely separate. Central anteriors nearer together than to the laterals. Anterior row slightly procurved or straight. Protarsus i. with a single small central apical spine beneath.

(Carapace precipitous behind. Labium rather over one-third the length of the maxillæ. Maxillæ longer, concave (or in species 4 and 5 not so concave) on the outer side. Trochanter of pedipalp attached distinctly below the middle of the maxilla.)

1. Anterior row of eyes slightly procurved. Maxillæ longer and straighter.

Legs variegated, black and orange. Abdomen with a pair of pale dorsal spots . . . . . *variegata*, sp. n.

2. Anterior row of eyes straight. Maxillæ shorter and rounded. Legs

orange-brown. Abdomen unicolorous, with a small semilunar white spot above the spinners . . . . . *semiluna*, sp. n.

NOTE.—The following characters apply to all the species falling under A in the above table; some, however, in the case of the eyes and the mouth-parts, apply also to B a and B b 2:—Eyes closely grouped; anterior row more or less procurved; eyes equidistant, or almost so. Labium not less than half the length of the maxillæ. Maxillæ short, broad, not straight, but curving inwards, convex on the outer side, not dilate at the apex. Trochanter of pedipalp attached more nearly at the middle of the maxilla on the outer side. No central apical protarsal spine. Carapace gradually inclined to its base, not precipitous behind the cephalic region, slightly convex at the central fovea, not concave, narrowed in front. Sternum scarcely longer than broad, almost circular. Lower margin of fang-groove with from 5–7 teeth.

### 1. *Corinna nigricans*.

*Corinna nigricans*, C. Koch, Die Arachniden, ix. p. 19, fig. 703<sup>1</sup>.

Type, ♂, in Mus. Berol. Total length 8 millim.

*Hab.* MEXICO<sup>1</sup>.

### 2. *Corinna raptor*.

*Hypsinotus raptor*, L. Koch, Die Drassiden, p. 274, t. 11. figg. 174, 175 (1866)<sup>1</sup>.

Type, ♂, in coll. Dr. L. Koch, Nürnberg. Length of carapace 6 millim.

*Hab.* MEXICO<sup>1</sup>.

This species, judging from Dr. Koch's figures and description, is quite different from any of those described below.

### 3. *Corinna spinifer*.

*Hypsinotus spinifer*, Keyserling, Verh. zool.-bot. Ges. Wien, xxxvii. p. 449, t. 6. fig. 20 (1887)<sup>1</sup>.

Type, ♀, in coll. Mus. Cambridge (Mass.).

*Hab.* NICARAGUA<sup>1</sup>.