

*Males.*

## A. Abdomen elongate, narrow.

1. Uncus of palpus almost straight from base to apex, the latter broader and irregularly serrate on the inner margin . . . . . *fuscovittata*, Keyserl.
2. Uncus of palpus strongly curved from base to apex, not serrate on inner margin, slightly recurved and aculeate at apex . . . . . *illicita*, O. P.-Cambr.

## B. Abdomen not elongate, triangular-oval.

- a. Conductor of palpus, lying beneath the apex of the embolus, much more prolonged, and more densely clothed with short setoid cusps. Apical lobes much more densely setose . . . . . *anastera*, Walck.
- b. Conductor of palpus quite short, very sparsely clothed, chiefly at the apex, with setoid cusps. Apical lobes very sparsely setose.
  - a<sup>1</sup>. Uncus of palpus with a strong transverse conical basal enlargement . . . . . *bifida*, sp. n.
  - b<sup>1</sup>. Uncus of palpus plain, without transverse enlargement at the base.
    - a<sup>2</sup>. Embolus of palpal bulb longer in proportion. Basal portion of uncus much broader, apex slightly recurved . . . . . *scutigera*, O. P.-Cambr.
    - b<sup>2</sup>. Embolus of palpal bulb shorter in proportion. Basal portion of uncus narrower, apex regularly curved, falciform.
  - a<sup>3</sup>. Uncus of palpus with an enlarged shoulder on the inner margin . . . . . *guttata*, sp. n.
  - b<sup>3</sup>. Uncus of palpus without enlarged shoulder on the inner margin . . . . . *vegeta*, Keyserl.

*Females.*

## A. Abdomen narrow, elongate.

1. Abdomen more prolonged behind, forming a sharp conical point; vulva much broader, not abruptly constricted near the apex . . . *fuscovittata*.
2. Abdomen less prolonged behind, forming a short, bluntly rounded cone; vulva narrower, abruptly constricted near its apex, then dilate . . . . . *illicita*.

## B. Abdomen broad, rounded-triangular.

- a. Ventral area of abdomen with a single central elongate-oval, or linear, white spot, immediately behind the genital rima.
  - a<sup>1</sup>. Hairs on the abdomen not enlarged basally, simply and gradually aculeate towards apex.
  - a<sup>2</sup>. Abdomen with a large central dorsal folium, with dentated margins. Scapus of vulva much longer.
  - a<sup>3</sup>. Abdomen not bituberculate at the apex; simply rounded, sometimes slightly conical.
  - a<sup>4</sup>. Scapus of vulva longer, broad basally, gradually or abruptly narrowed to a fine point; the basal portion transversely and strongly striate.
  - a<sup>5</sup>. Scapus not abruptly narrowed . . . . . *anastera*.