

26. **Aranea graphica.** (Tab. LI. figg. 2, 2 a, ♂.)

Epeira graphica, O. P.-Cambr. Biol. Centr.-Amer., Arachn. Aran. i. p. 22, t. 7. figg. 16, 16 a-c (♂)¹; Keyserl. Spinn. Amer., Epeiridae, p. 121, t. 6. figg. 89, 89 a (♂)².

Type ♂, in coll. Godman & Salvin. Total length 3.5 millim.

Carapace yellow, cephalic area brown; legs pale yellow-white, suffused with dusky brown; mandibles pale yellow; sternum pale yellow, with a blackish central spot. Abdomen mottled with black and pale yellow, with a wide central folium margined with black spots, and a conspicuous anterior, dorsal, central, elongate white spot. Femora and protarsi i. and ii. without spines beneath. Tibia ii. not incrassate, with 4—2 long spines on the inner side. Patella of pedipalp with a single spine at the apex, that of the legs without apical spine (the scars only appear on the apex of the patella of the pedipalp). The embolus of the palpal bulb is quite characteristic, see Plate.

Hab. GUATEMALA²; PANAMA, Bugaba (*Champion*¹).

27. **Aranea flava.** (Tab. LI. fig. 3, ♂.)

Singa flava, O. P.-Cambr. Biol. Centr.-Amer., Arachn. Aran. i. pp. x, 136¹.

Singa lutea, O. P.-Cambr. loc. cit. t. 17. figg. 15, 15 a-d (♂)².

Type, ♂, in coll. Godman & Salvin. Total length 3 millim.

Hab. MEXICO, Tierra Colorada (*H. H. Smith*¹).

28. **Aranea cyrtophoroides**, sp. n. (Tab. LI. figg. 4, 4 a-d, ♀.)

Type, ♀, in coll. Godman & Salvin. Total length 10 millim.

Carapace, legs, sternum, and mandibles dull orange-brown. Abdomen clay-yellow-brown, with a shoulder-cusp on each side, anteriorly; ventral area dark, with a narrow curved white band on each side. Both rows of eyes slightly recurved; central quadrangle longer than broad, very slightly narrower behind, eyes subequal, posteriors one diameter, anteriors slightly more than a diameter, apart; laterals well separate, one diameter apart. Legs without stout spines, but with fine bristles in their place. Femora with a row of fine bristles on each side beneath; tibiae and protarsi with a double row of bristles beneath, those on the protarsi being stouter; patellae with a fine apical bristle. Mandibles with three teeth on the lower margin. Sternum as broad as long. Abdomen somewhat prolonged beyond the spinners above; the dorsal muscular scars very conspicuous. For the vulva, see Plate.

Hab. MEXICO, Teapa (*H. H. Smith*).

29. **Aranea ocellata.**

Epeira ocellata, O. P.-Cambr. Biol. Centr.-Amer., Arachn. Aran. i. p. 29, t. 6. figg. 17, 17 a (♀)¹.

Type, ♀ (immature), in coll. Godman & Salvin. Total length 4.5 millim.

Hab. GUATEMALA, Cahabon (*Sarg*¹).

30. **Aranea consequa.**

Epeira consequa, O. P.-Cambr. Biol. Centr.-Amer., Arachn. Aran. i. pp. x, 36 (♀)¹.

Epeira destriata, O. P.-Cambr. loc. cit. t. 14. figg. 14, 14 a, b (♀, nec ♂)².

Type, ♀, in coll. Godman & Salvin. Total length 8 millim.

Hab. MEXICO, Ciudad in Durango (*Forrer*¹); PANAMA, Bugaba (*Champion*¹).

The types of this species have been mislaid.