

31. *Aranea conica*.

Alpaida conica, O. P.-Cambr. Biol. Centr. Amer., Arachn. Aran. i. p. 53, t. 4. figg. 3, 3 *a-e* (♀)¹;
Keyserl. Spinn. Amer., Epeiridæ, p. 281, t. 14. figg. 207, 207 *a, b* (♀)².

Type, ♀, in coll. Godman & Salvin. Total length 5 millim.

Hab. GUATEMALA²; PANAMA, Bugaba (*Champion*¹).

32. *Aranea sallei*.

Epeira salei (sic), Keyserl. Sitzungsab. nat. Ges. Isis, Dresden, 1863, p. 93, t. 4. figg. 10, 11 (♀)¹;
Spinn. Amer., Epeiridæ, p. 196, t. 9. figg. 145, 145 *a* (♀)².

Type, ♀, in Mus. Brit. Total length 8.5 millim.

Hab. MEXICO, Oaxaca (*Sallé*^{1 2}).

33. *Aranea rivalis*.

Epeira rivalis, Keyserl. Spinn. Amer., Epeiridæ, p. 103, t. 5. figg. 76, 76 *a* (♀)¹.

Type, ♀, in coll. Marx. Total length 12.2 millim.

Hab. GUATEMALA¹.

34. *Aranea nicaraguensis*.

Epeira nicaraguensis, Keyserl. Verh. zool.-bot. Ges. Wien, xxxiv. p. 532, t. 13. fig. 31 (♀)
(1884)¹; Spinn. Amer., Epeiridæ, p. 214, t. 10. figg. 159, 159 *a* (♀)².

Type, ♀, in Mus. Cambridge, U.S.A. Total length 17 millim.

Hab. NICARAGUA^{1 2}.

35. *Aranea zilloides*.

Epeira zilloides, Banks, Proc. Calif. Acad. Sci. (3) i. p. 255, t. 15. fig. 2 (♂ ♀)¹.

Type ♂, gynetype ♀, in coll. Calif. Acad. Sciences. Total length, ♂ 4, ♀ 7.5 millim.

Hab. MEXICO, Tepic¹.

36. *Aranea fasciolata*. (Tab. LI. figg. 5, 5 *a*, ♂.)

Epeiroides fasciolata, O. P.-Cambr. Biol. Centr.-Amer., Arachn. Aran. i. p. 15, t. 8. figg. 5,
5 *a-e* (♂)¹.

[Nec Keyserl. Spinn. Amer., Epeiridæ, p. 309, t. 16. figg. 228, 228 *a-c* (♂).]

Type, ♂, in coll. Godman & Salvin. Total length 2 millim.

Hab. PANAMA, Bugaba (*Champion*¹).

This species probably belongs to the group Theridiosommatinæ. The spider figured and described under this name by Keyserling, which is not in the British Museum, was perhaps incorrectly identified, as no long bristles on the first pair of legs are represented on his plate.