

- a*<sup>4</sup>. Posterior spinning-mammillæ very long, flagelliform, tapering, attenuate . . . . . HERSILIIDÆ (p. 350).
- b*<sup>4</sup>. Posterior spinning-mammillæ not abnormally long or flagelliform.
- \* Tarsal claws 2. Ungual tufts present.
- a*<sup>5</sup>. Anterior pair of spinning-mammillæ set wide apart at their base . . . . . GNAPHOSIDÆ (p. 54).
- b*<sup>5</sup>. Anterior pair of spinning-mammillæ set close together at their base.
- a*<sup>6</sup>. Apex of protarsi without a hard chitinous rim, but furnished with a trilobate membrane . . . . . HETEROPODIDÆ (p. 118).
- b*<sup>6</sup>. Apex of protarsi with a hard chitinous rim, not furnished with a trilobate membrane.
- a*<sup>7</sup>. Eyes situated in an almost straight transverse row . . . . . SELENOPIDÆ (p. 115).
- b*<sup>7</sup>. Eyes situated in two or three transverse rows.
- a*<sup>8</sup>. Legs iii. and iv. very much shorter than i. and ii. Mandibles weak, sometimes with very small teeth . . . . . THOMISIDÆ (p. 127).
- b*<sup>8</sup>. Legs iii. and iv. not much shorter than i. and ii. Mandibles strong, with stout teeth.
- a*<sup>9</sup>. Central anterior eyes very large. Eyes of posterior row widely separate, forming a large quadrangle . . . . . SALTICIDÆ (p. 166).
- b*<sup>9</sup>. Central anterior eyes not or scarcely enlarged. Eyes of posterior row not widely separate, forming a transverse row.
- a*<sup>10</sup>. Eyes situated in two transverse rows.
- a*<sup>11</sup>. Tracheal stigmata more or less remote from the spinning-mammillæ . . . . . ANYPHÆNIDÆ (p. 92).
- b*<sup>11</sup>. Tracheal stigmata set immediately in front of the spinning-mammillæ . . . . . CLUBIONIDÆ (p. 61).
- b*<sup>10</sup>. Eyes situated in three, or four, transverse rows . . . . . CTENIDÆ (p. 107).
- \*\* Tarsal claws 3.
- a*<sup>12</sup>. Tibia of male palpus with an apical external spur.
- a*<sup>13</sup>. Protarsi i. and ii. without a central apical spine beneath . . . . . SENOCULIDÆ (p. 349).