

- a*<sup>13</sup>. Femur iv. with two series, a dorsal and an inferior, of comparatively short denticles. No external basal series.
- a*<sup>14</sup>. The inferior series of denticles on femur iv. subequal in length throughout; the basal denticle erect and set out of the plane of the rest of the series . . . *erectispinus*, sp. n.
- b*<sup>14</sup>. The inferior series of denticles on femur iv. becoming much longer from the middle towards the apex; the basal denticle quite small and normally situated . . . *rectus*, sp. n.
- b*<sup>13</sup>. Femur iv. with three series of very long denticles, those of the inferior series as long as three times the width of the segment; also with a short external basal series of four long denticles . . . *pectinigerus*, sp. n.

*Females.*

A. Tarsus iii. with 7 segments.

- a*. Third segment of dorsal scutum with a pair of rudimentary tubercles only. Coxa iv. with a double tubercle at its base, just behind the trochanter iii. Size much longer, length 9 millim. . . . *serrato-tibialis*.
- b*. Third segment of dorsal scutum with a pair of sharp spines. Coxa iv. without tubercles as above. Size much smaller, length 5 millim.
- a*<sup>1</sup>. Femur iv. longer (5 millim.), and scarcely arcuate in the basal half . . . *simplicipes*.
- b*<sup>1</sup>. Femur iv. shorter (4 millim.), more strongly arcuate in the basal half . . . *sinuosus*.

B. Tarsus iii. with 8 segments.

- a*<sup>2</sup>. Tarsus iv. with 8 segments . . . *serratipes*.
- b*<sup>2</sup>. Tarsus iv. with 9 segments.
- a*<sup>3</sup>. Size much larger, length 9 millim. Third segment of dorsal scutum entirely devoid of spines or tubercles. Femur iv. arcuate in the basal half . . . *pectinigerus*.
- b*<sup>3</sup>. Size much smaller, length 5 millim. Third segment of dorsal scutum with a pair of long stout spines. Femur iv. straight throughout . . . *erectispinus*.

1. *Erginus serratifer*, sp. n. (Tab. LII. fig. 18, ♂.)

Type, ♂, in coll. Godman & Salvin. Total length 5 millim. (excl. mand.); femur iv. 3.25 millim.

- ♂. General characters similar to those of *E. pectinigerus*. The two spines on the tergal third segment are short. Tarsal segments, i. 6; ii. 13; iii. 7; iv. 7. Trochanters iii. and iv. and coxa iv. with an apical tooth. Basal segment of mandible granulose and in other respects similar to that of *E. erectispinus*. Femur iv. with an internal inferior series of 18 long denticles, basally coalescent, the longest more than twice the width of the segment, extending from the base, but not quite reaching the apex, and an external, dorsal, basal series of 5 or 6 shorter denticles occupying the basal third, besides two inferior series of small tubercles extending from the base, but not quite reaching the apex, those of the