

7. *Trombidium albicolle*, sp. n. (Tab. I. figg. 3, 3 a.)

Corpus oblongum, humeris protractis, apice anali rotundato; coccineum, hirsutulum, maculis atque striis albis indutum; in apice frontali pseudo-prothoracis macula alba; stria transversa lata inter humeros, postice linguam latam medianam in abdominis dorsum emittens; apex analis albus; duæ maculæ albæ latero-marginales ante corporis apicem sitæ, parvæ. Coxæ lateribus corporis infixæ. Pedes longi, graciles, rufescentes, inter se fere æquales. Palpi graciles, pilis longis rarioribus instructi; tertio articulo elongato, cylindrico; quarti ungue unidentato, appendicula recta, apice rotundata, ad apicem piligera.

Long. 0·75–1 millim.

Hab. GUATEMALA, Antigua (*Stoll*).

Body oblong, rounded behind, the shoulders protruding; pseudo-prothorax triangular: colour scarlet, with white spots formed by white, thickly-set, velvety hairs; these spots are arranged as follows—a round white one on the frontal surface of the pseudo-prothorax, a large T-shaped one on the middle of the dorsal surface behind the eyes, an anal one at the end of the body, and two small round lateral ones near the margins of the posterior part of the abdomen. Legs, palpi, and mandibles of a clear reddish colour. Legs long, slender, the first and fourth pairs a trifle longer than the second and third. Third joint of the palpi long, cylindrical; the fourth joint with a small tooth on the concave side of its claw, the appendicula straight, parallel-sided, rounded at the end, where some stiff hairs are inserted.

This species is found with the preceding, on bushes in hedges and gardens in the valley of Antigua.

8. *Trombidium muricola*, sp. n. (Tab. II. figg. 3–3 b.)

Corpus breve, obovoideum, convexum, humeris rotundatis, apice frontali rotundata; cuticula lævis, sericeo-nitida, nigra, maculis et striis albis, variantibus figura. Pedes, palpi et mandibulæ rufescentes. Pedes longi, graciles; primum par cæteris longius.

Hab. GUATEMALA, Antigua (*Stoll*).

Body short, ovoid, convex, with the shoulders and the frontal and anal apex rounded; smooth, silky-black, with white spots and stripes varying in form in the different specimens: in some the dorsal surface of the abdomen bears a white triangular spot on its posterior third, and from the acute angle of this, which is directed forwards, two narrow stripes proceed obliquely towards the humeri; in others there only remains a small white spot at the anal end, and a narrow white streak running across the dorsum at a short distance behind the middle, the streak in its centre forming a large angle (sometimes connected with the anal spot by a narrow white line) which opens towards the frontal apex; finally, there occur specimens in which the whole body is black, except a small white border along the posterior margin. The legs are long and slender, the front pair a little longer than the others. Legs, palpi, and mandibles light reddish-brown.

This species is found in the rainy season on the adobe-walls of the nopal-gardens