

dorsally, but never replaced by a secondary sulcus lying in front of the pore; phallopod, where known, with its distal segment long and slender, the guard of the seminal style much shorter than the segment.

- b.* Scobina unspecialized, the extremities of the groove defining the impression turned outwards, *i. e.* away from each other; the striæ of the striate area very coarse, hardly more numerous and hardly finer than those of the adjacent area of the terga; the longitudinal striæ of the posterior portion of the segment extending at least as high as the pores. *olivaceus*, Newp.
- b*<sup>1</sup>. Scobina specialized, the extremities of the groove incurled towards each other, thus forming a crescentic or subcrescentic impression; the striæ of the striate area very fine and numerous; longitudinal grooves of posterior portion of segments not extending halfway up to the pore.
- c.* Sensory papillæ on apical segment of antennæ very numerous, variable in number, and close-set.
- d.* Scobina very small and not extending posteriorly as far as the middle segment of the body; infero-lateral edge of first tergal plate not widely rounded, subangular, its anterior border slightly cut away. *rogersi*, sp. n.
- d*<sup>1</sup>. Scobina large, extending posteriorly past the middle segment of the body; infero-lateral edge of first tergal plate evenly rounded.
- e.* Sternal plate of anal segment very short, longitudinally convex, and not or hardly longer than the sternal area of the preceding segment, its posterior border straight and transverse; dorsal area of the segments irregularly tubercular in front. *marci*, sp. n.
- e*<sup>1</sup>. Sternal plate of anal segment at least twice as long as the sternal area of the preceding segment, its posterior border angular or strongly convex; no tubercles on dorsal area of the segments.
- f.* Antennæ much longer than head; legs and antennæ dark, at least in their basal half. *rixi*, sp. n.
- f*<sup>1</sup>. Antennæ shorter or at least not longer than head; legs and antennæ uniformly pale.
- g.* Colour crimson; sternal plate of anal segment with convex posterior edge, and marked in front with a transverse groove. *aposematus*, sp. n.
- g*<sup>1</sup>. Colour red, banded with green (? black); sternal plate of anal segment with angular edge and no anterior groove. *tristani*, sp. n.
- c*<sup>1</sup>. Sensory papillæ on apical segment of antennæ few in number (4-5) and widely spaced.
- h.* Scobinæ very wide, each about twice as wide as the area between them on the middle segments of the body; inferior horizontal area of second segment with a deep transverse groove behind the thickened anterior edge. *scobinatus*, sp. n.
- h*<sup>1</sup>. Scobinæ narrower, separated by a space which is not less than twice the transverse diameter of each; inferior horizontal area of second segment shorter, without any transverse groove behind the thin upstanding anterior edge.