

OZÆNA.

Ozæna, Olivier, Encycl. Méth. viii. pt. 2, p. 617 (1791); Chaudoir, Ann. Soc. Ent. Belg. xi. p. 50 (1818).

Ictinus, Castelnau, Etud. Ent. p. 53 (1835).

Also peculiar to Tropical America; three species have been described.

1. *Ozæna brevicornis*.

Ozæna brevicornis, Bates, Ent. Monthly Mag. xi. p. 24¹.

Hab. NICARAGUA, Chontales (*Belt*).—SOUTH AMERICA, Peru¹?

PACHYTELES.

Pachyteles, Perty, Del. An. Art. Brasil. p. 3, t. i. f. 10 (1830).

Ozæna (partim), Dejean, Sp. Gén. i. p. 433.

Ictinus (partim), Castelnau, Etud. Ent. p. 145 (1835).

Goniotropis, Gray, Griffith's An. King. i. p. 273 (1832).

Tropopsis, Solier, Gay, Hist. Chile, Zool. iv. p. 179 (1849).

Upwards of sixty species of this genus are known by descriptions. Two of the number are from West Tropical Africa, and one from Madagascar; the rest are exclusively American, ranging from Arizona in the north to Monte Video and Chili in the south.

1. *Pachyteles porrectus*.

Pachyteles porrectus, Chaudoir, Ann. Soc. Ent. Belg. xi. p. 54¹ (1868).

Hab. MEXICO, Puebla (*Boucard*¹), Playa Vicente (*Sallé*); BRITISH HONDURAS (*Blancaneaux*); GUATEMALA, Izabal (*Sallé*), Panzos, San Isidro, Pantaleon (*Champion*); NICARAGUA, Chontales (*Belt*).

2. *Pachyteles funcki*. (Tab. II. fig. 5.)

Pachyteles Funckii, Chaudoir, Ann. Soc. Ent. Belg. xi. p. 55 (1868)¹.

Hab. NICARAGUA, Chontales (*Belt*, *Janson*).—SOUTH AMERICA, Venezuela¹.

3. *Pachyteles verticalis*. (Tab. II. fig. 16.)

Ozæna verticalis, Chaudoir, Bull. Mosc. 1848, i. p. 104¹.

Pachyteles verticalis, Chaudoir, Ann. Soc. Ent. Belg. xi. p. 56 (1868)².

Hab. MEXICO¹, Cordova (*Sallé*²), Jalapa, Las Vigas (*Höge*); GUATEMALA, Zapote, Pantaleon, Las Mercedes (*Champion*); NICARAGUA, Chontales (*Janson*).

In some examples, chiefly those of Nicaragua and Guatemala, the anterior and
BIOL. CENTR.-AMER., Coleopt., Vol. I. Pt. 1, December 1881. ee