

CLIVINA.

Clivina, Latreille, Considérations Générales s. l. Crust. Arachn. et Ins. p. 156 (1810); Putzeys, Mém. Ac. R. Liège, ii. 1846, p. 59.

Similar in its wide distribution to *Dyschirius*, and even more numerous in species; a larger proportion, however, have been described from the tropical zone, and it is well represented in Central and other parts of Tropical America. About 220 species are known.

1. *Clivina planulata*.

Clivina planulata, Putzeys, Révis. Gén. p. 140¹.

Hab. MEXICO (*Deyrolle*¹).

2. *Clivina planicollis*.

Clivina planicollis, Leconte, Proc. Acad. Phil. 1857, p. 81¹; Putzeys, Révis. Gén. p. 142².

Hab. NORTH AMERICA, United States¹.—MEXICO², Vera Cruz, Cordova, La Soledad (*Sallé*).

3. *Clivina lucida*.

Clivina lucida, Putzeys, Révis. Gén. p. 150¹.

Hab. MEXICO (*Riehl*¹).

4. *Clivina distigma*. (Tab. II. fig. 24.)

Clivina distigma, Putzeys, Révis. Gén. p. 151¹.

Hab. MEXICO¹, Cordova (*Sallé*), Valle del Maiz (*Dr. Palmer*), Jalapa (*Höge*); GUATEMALA, San Gerónimo (*Champion*).

5. *Clivina cruciata*.

Clivina cruciata, Putzeys, Révis. Gén. p. 152.

Hab. MEXICO, Tuxtla (*Sallé*); GUATEMALA, Zapote (*Champion*).

6. *Clivina brunneipennis*.

Clivina brunneipennis, Putz. Monogr. Clivinides, p. 107¹.

Clivina brunneipennis, id. Révis. Gén. p. 159.

Hab. MEXICO¹, San Luis Potosi (*Dr. Palmer*), Oaxaca (*Höge*); GUATEMALA, San Gerónimo (*Champion*).

7. *Clivina bituberculata*.

Clivina tuberculata, Putzeys, Révis. Gén. p. 161¹.

Hab. MEXICO¹.