

2. *Pelecium scydmænoides*.

Eripus scydmænoides, Dejean, Sp. Gén. Col. iv. p. 10¹.

Pelecium scydmænoides, Chaudoir, Berl. entom. Zeitschr. 1861, p. 129.

Hab. MEXICO¹.

3. *Pelecium suturale*.

Pelecium suturale, Chaudoir, Berl. entom. Zeitschr. 1861, p. 129¹.

Hab. MEXICO¹, Cordova (*Sallé*).

4. *Pelecium microphthalmum*.

Pelecium microphthalmum, Chaudoir, Rev. et Mag. Zool. 1866, p. 108¹.

Hab. MEXICO¹, Orizaba (*Sallé*).

5. *Pelecium globipenne*.

Pelecium globipenne, Chaudoir, Rev. et Mag. Zool. 1866, p. 109¹.

Hab. MEXICO¹, Orizaba (*Sallé*).

6. *Pelecium subdentatum*.

Pelecium subdentatum, Chaudoir, Rev. et Mag. Zool. 1866, p. 109¹.

Hab. MEXICO¹.

7. *Pelecium subcæcum*.

Pelecium subcæcum, Chaudoir, Rev. et Mag. Zool. 1866, p. 110¹.

Hab. MEXICO¹.

Subdivision B. *Patellimani* (Latreille, Dejean).

Subfamily *PANAGÆINÆ*.

PANAGÆUS.

Panagæus, Latreille, Hist. Nat. Crust. et Ins. viii. p. 291 (1804); Chaudoir, Essai Monogr. sur les Panagéides, p. 92 (1878).

As restricted by Chaudoir this genus contains only nine known species, and is confined to the northern hemisphere in both the new and old worlds.

1. *Panagæus panamensis*.

Panagæus panamensis, Laferté, Ann. Soc. Ent. Fr. 1851, p. 223; Chaudoir, Bull. Mosc. 1861, iv. p. 352¹; id. Essai Monogr. sur les Panagéides, p. 96.

Hab. PANAMA¹.