

2. *Panagæus quadrisignatus*.

Panagæus quadrisignatus, Chevrolat, Col. Mex. ii. no. 187 (1835)¹; Chaudoir, Essai Monogr. s. l. Panagéides, p. 97².

Panagæus Thomæ, Schaum, Ann. Soc. Ent. Fr. 1853, p. 441³.

Panagæus mexicanus, Putzeys, Mém. Soc. Sc. Liège, ii. 1846, p. 401.

Hab. MEXICO², Vera Cruz (*Sallé*)¹.—WEST INDIES, St. Thomas^{2 3}.

3. *Panagæus sallæi*. (Tab. III. fig. 2.)

Panagæus Sallei, Chaudoir, Bull. Mosc. 1861, ii. p. 353; id. Essai Monogr. p. 99.

Hab. MEXICO, Puebla, Yolos, Aculsingo (*Sallé*), Jalapa, Tehuacan (*Höge*).

Subfam. *CHLÆNIINÆ*.

CHLÆNIUS.

Chlænius, Bonelli, Observations Entomologiques, i. tabl. synopt. (1809); Chaudoir, Monographie des Chlénien, p. 10 (1876).

In his recent monograph of this numerous and beautiful genus, the late Baron Chaudoir enumerated 404 species as belonging to it. It is nearly restricted to continental lands and islands, the few species found in suboceanic islands, such as the Canaries, New Caledonia, and New Guinea, being common insects in the nearest large tract of land. In America the genus is well represented on the northern continent and in the basin of the La Plata; but the number of species which occur in the equatorial parts of the continent is very limited.

1. *Chlænius soginoides*.

Chlænius soginoides, Chaudoir, Monogr. Chlén. p. 74¹.

Hab. MEXICO, Guanajuato¹, city of Mexico (*Flohr*, *Dr. Palmer*).

2. *Chlænius validus*.

Omaseus validus, Chevrolat, Col. Mex. ii. no. 174¹.

Chlænius (*Glyptoderus*) *validus*, Chaudoir, Monogr. Chlén. p. 75².

Hab. MEXICO^{1 2}, Mirador (*Sallé*), Jalapa (*Höge*).

3. *Chlænius chrysopleurus*.

Chlænius chrysopleurus, Chaudoir, Bull. Mosc. 1856, ii. p. 276.

Chlænius (*Glyptoderus*) *chrysopleurus*, Chaudoir, Monogr. Chlén. p. 78¹.

Glyptoderus aurolimbatus, Laferté, Ann. Soc. Ent. Fr. 1851, p. 261.

Hab. MEXICO, Yucatan¹, Campeche (*Sallé*); NICARAGUA, Chontales (*Belt*, *Janson*).

Var. *cærulea*, Chaud. Monogr. p. 78¹ (*amabilis*, Dugès in coll. *Sallé*²).

Hab. Guanajuato² (*Dugès*), Puebla¹ (*Sallé*).

Var. thoracis plerumque ut in typo disco cupreo, sed elytris marginibus anguste viridi-cæruleis.

Hab. MEXICO, Oaxaca (*Höge*); GUATEMALA (*Sallé*), near the city (*Salvin*).

BIOL. CENTR.-AMER., Coleopt., Vol. I. Pt. 1, February 1882.

gg