

4. *Chlænius menevillei*. (Tab. III. fig. 3.)

Chlænius (*Glyptoderus*) *Menevillei*, Chaudoir, Monogr. Chlén. p. 77¹.

Glyptoderus Guerini, Ann. Soc. Ent. Fr. 1851, p. 261 (nom. præocc.).

Ch. chrysopleuro proxime affinis, sed differt elytris grossius et sparsius punctulatis striisque valde crenato-punctatis. Nitidior, niger, capite, thorace elytrisque marginibus læte viridi-æneis, thoracis disco nigro-cupreo nitido, latera versus splendide cupreo; elytris fortiter striatis, striis grosse transversim punctatis, interstitiis convexis nitidis subsparsim setifero-punctatis.

Long. 8 lin. ♂ ♀.

Hab. PANAMA (*Boucard*), Volcan de Chiriqui (*Champion*).—SOUTH AMERICA, Bolivia¹.

A specimen communicated by M. René Oberthür, doubtless compared with Chaudoir's type, was labelled as from Panama. The locality Bolivia is probably erroneous. The specimen figured is from Chiriqui.

5. *Chlænius cæruleicollis*.

Chlænius cæruleicollis, Chaudoir, Monogr. Chlén. p. 78¹.

Hab. MEXICO, Las Peras¹, Capulalpam, Puebla, Toluca (*Sallé*), Jalapa, La Parada (*Höge*).

6. *Chlænius aberrans*. (*Chlænius martinezi*, Tab. III. fig. 4.)

Ch. cæruleicollis multo major et magis elongatus, elytris elongato-ovatis, thorace relative parvo, cordato-quadrato, antice valde rotundato, postice sinuatim modice angustato angulis posticis rectis; elytris sulculo marginali apud humeros angulato; episternis prothoracis rugulosis, metathoracis grosse punctatis extus haud sulcatis; capite viridi-æneo, medio lævi, lateribus punctatis, collo depresso violaceo; thorace grosse discrete subrugulose punctato, violaceo, lateribus viridi-æneis; elytris acute sed haud profunde striatis, interstitiis subtiliter sparsim punctulatis, opacis, violaceis, limbo laterali viridi opaco; antennis orisque partibus piceo-rufis, pedibus nigro-piceis.

Long. 8½ lin. ♂.

Hab. MEXICO (*Sallé*).

I have adopted the MS. name given to this species in the Chaudoir collection, now in the possession of M. René Oberthür. Chaudoir gave the name *Martinezi* to the same species in the Sallé collection.

7. *Chlænius scabricollis*.

Chlænius scabricollis, Chevrolat, Col. Mex. i. fasc. 2 (1833); Chaudoir, Monogr. Chlén. p. 96.

Hab. MEXICO, Tuxtla (*Sallé*).

8. *Chlænius sallæi*. (Tab. III. fig. 5.)

Chlænius sallæi, Chaudoir, Monogr. Chlén. p. 149.

Hab. MEXICO, Cordova (*Sallé*).

9. *Chlænius rodriguezi*.

Chlænius rodriguezi, Chaudoir, Monogr. Chlén. p. 152¹.

Hab. GUATEMALA, Coban¹ (*Sallé*), Capetillo (*Champion*).

10. *Chlænius leucoscelis*.

Chlænius leucoscelis, Chevrolat, Col. Mex. i. fasc. 4 (1834)¹; Chaudoir, Monogr. Chlén. p. 190.

Hab. MEXICO¹, Cordova, Izucar, Etla, Capulalpam, Guanajuato (*Sallé*), Oaxaca (*Höge*); GUATEMALA, near the city (*Champion*).