

11. **Chlænius chlorochrous.**

Chlænius chlorochrous, Chaudoir, Monogr. Chlén. p. 190¹.

Hab. MEXICO¹, Playa Vicente (*Sallé*); GUATEMALA, Rio Maria Linda, Panajachel (*Champion*).

12. **Chlænius ruficauda.** (Tab. III. fig. 7.)

Chlænius ruficauda, Chaudoir, Bull. Mosc. 1856, ii. p. 194¹.

Chlænius apicalis, Leconte, Ann. Lyc. N. York, v. p. 179 (1851)², nom. præocc.

Chlænius posticus, Leconte, Journ. Acad. Phil. iv. 1858, t. 1. f. 8².

Chlænius dimidiatus, Motschulsky, Etud. Ent. 1858, p. 157.

Hab. NORTH AMERICA, California².—MEXICO¹, Puebla (*Sallé*), Juquila, Oaxaca (*Höge*).

13. **Chlænius herbaceus.**

Chlænius herbaceus, Chevrolat, Col. Mex. ii. no. 188 (1835)¹; Chaudoir, Monogr. Chlén. p. 212¹.

Chlænius patruelis, Leconte, Ann. Lyc. N. York, iv. p. 435 (1848)².

Hab. NORTH AMERICA, Georgia².—MEXICO¹, Guanajuato (*Sallé*).

14. **Chlænius guatemalenus.** (Tab. III. fig. 6.)

Ch. cursori affinissimus, sed differt elytrorum interstitiis subgrosse punctatis. Supra viridi-æneus vel violaceus, thorace medio elytrisque versus suturam nigricantibus; antennis, partibus oris pedibusque flavo-testaceis; capite (media fronte excepta) et thorace grosse sparsim punctatis, hoc quadrato-cordato, prope angulos anticos angustato, versus basin sinuato, angulis posticis oblique truncatis; elytris elongato-ovalibus margine basali flexuoso, humeris angulatis, mox pone humeros ampliatis, profunde striatis striis subtiliter punctulatis, interstitiis grosse sparsim punctatis; abdomine utrinque modice punctulato.

Long. $7\frac{1}{2}$ lin.

Hab. GUATEMALA, Paso Antonio (*Champion*).

Differs from *C. cursor* too widely in the punctuation of the elytral interstices to be referred to that species as a variety. *C. cursor* varies much in the convexity of the interstices of the elytra, M. Sallé's named examples and all those taken by Höge at Jalapa and Oaxaca having them distinctly convex. The species is also allied to *C. rodriguezi*, which is placed by Chaudoir in a different primary division of the genus, according to his complicated classification.

15. **Chlænius cursor.**

Chlænius cursor, Chevrolat, Col. Mex. ii. no. 168 (1835)¹; Chaudoir, Monogr. Chlén. p. 212².

Chlænius sparsus, Leconte, List of Col. New Species, 1863, p. 12³.

Hab. NORTH AMERICA, Lower California³.—MEXICO, Orizaba¹, Cordova, Tuxtla, Panistlahuca (*Sallé*), Guanajuato², Jalapa, Oaxaca (*Höge*).

16. **Chlænius orbus.**

Chlænius orbus, Horn, Trans. Am. Ent. Soc. 1871, p. 326¹; Chaudoir, Monogr. Chlén. p. 216.

Hab. NORTH AMERICA, Texas¹.—MEXICO, Oaxaca (*Höge*).