

17. *Chlænius azurescens*.

Chlænius azurescens, Chaudoir, Monogr. Chlén. p. 220¹.

Hab. MEXICO, Orizaba¹, Playa Vicente (*Sallé*), Almolonga (*Höge*).

18. *Chlænius planipennis*. (Tab. III. fig. 8.)

Chlænius planipennis, Chaudoir, Monogr. Chlén. p. 220.

Hab. MEXICO, Juquila (*Sallé*).

19. *Chlænius tricolor*.

Chlænius tricolor, Dejean, Spec. Gén. Col. ii. p. 334ⁿ; Chaudoir, Monogr. Chlén. p. 241.

Var. *Chlænius vigilans*, Say, Trans. Am. Phil. Soc. iv. p. 419 (1834)².

Chlænius chalybeipennis, Chevrolat, Col. Mex. i. fasc. 4 (1834)³.

Hab. NORTH AMERICA¹, northwards to Canada.—MEXICO^{1 2 3}, Guanajuato (*Dugès*, coll. *Sallé*), Oaxaca (*Sallé*, *Höge*), Almolonga (*Höge*); GUATEMALA (*Sallé*), Panajachel (*Champion*).

The species occurs in Mexico and Guatemala under two distinct colour varieties. In one, which prevails at Oaxaca and in Guatemala, the head and thorax are green; in the other, occurring chiefly near the capital and at Jalapa and Puebla, the same parts are of a rich coppery hue: some Jalapa specimens are intermediate in colour. The green variety is that described by Say as *Chlænius vigilans*.

20. *Chlænius melanarius*.

Chlænius melanarius, Dejean, Sp. Gén. Col. v. p. 658 (1831)¹; Chaudoir, Monogr. Chlén. p. 242.

Chlænius soccatus, Say, Trans. Am. Phil. Soc. iv. p. 419 (1834)².

Chlænius Chevrolati, Chevrolat, Col. Mex. ii. no. 171 (1835).

Var. *Chlænius violaceus*, Chevrolat, Col. Mex. ii. no. 170 (1835); Chaudoir, Monogr. Chlén. p. 242.

Hab. MEXICO^{1 2}, Guanajuato, Puebla, Toluca (*Sallé*), Hacienda de los Bleados (*Dr. Palmer*), Mexico city (*Sallé*, *Dr. Palmer*, *Höge*), Jalapa, Oaxaca (*Höge*).

The two forms (pale legs and black legs) occur together in the same localities.

21. *Chlænius obsoletus*.

Chlænius obsoletus, Leconte, Ann. Lyc. N. York, v. p. 180¹; Chaudoir, Monogr. Chlén. p. 244².

Hab. NORTH AMERICA, California¹.—MEXICO², Oaxaca (*Höge*); GUATEMALA (*Sallé*), Guatemala city (*Champion*); NICARAGUA (*Sallé*).

22. *Chlænius viridicollis*.

Chlænius viridicollis, Reiche, Rev. Zool. 1843, p. 37¹; Chaudoir, Monogr. Chlén. p. 244².

Hab. MEXICO², Etla (*Sallé*), Oaxaca (*Höge*).—SOUTH AMERICA, Colombia¹, Amazons².