

7. *Selenophorus seriatoporus*.

Selenophorus seriatoporus, Putzeys, Stett. ent. Zeit. 1878, p. 23¹.

Hab. MEXICO, Cordova¹, Yolos (*Sallé*), Oaxaca (*Höge*); GUATEMALA, near the city, San Gerónimo, El Reposo, Paso Antonio (*Champion*).

Guatemalan examples are very generally shorter in form and more bronzed in colour than those of Mexico. The difference is not sufficiently marked or constant for specific distinction or for a well-defined race.

8. *Selenophorus callistichus*.

Selenophorus callistichus, Bates, Proc. Zool. Soc. 1878, p. 591¹.

"Oblongo-ovatus, elongatus, minus convexus, cupreo-æneus; antennis basi pedibusque flavo-testaceis; thorace antice gradatim, versus basin citius, angustato, angulis posticis rotundatis, basi tota haud profunde punctulata; elytris apice sinuatis, supra fortiter striatis, interstitiis alternis punctis numerosis rotundatis seriatim impressis; tarsis posticis gracilibus, elongatis; tibiis et tarsis absque spinis fortibus.

"Long. $4\frac{1}{2}$ lin. ♂ ♀."

Hab. PANAMA, Obispo (*Salvin*¹), Volcan de Chiriqui, alt. 2000 to 3000 feet (*Champion*).

Distinguished from *S. seriatoporus*, particularly the typical Mexican form, by its richer metallic colour and the larger and more conspicuous punctures of the elytra.

9. *Selenophorus chiriquinus*.

S. seriatoporo et *callisticho* affinis, sed multo minor, lætius æneus, haud sericeo-nitens, thoraceque basi grossius punctato. Æneus, antennis basi pedibusque flavo-testaceis; thorace lateribus modice æqualiter arcuatis, tota basi subgrosse et crebre punctata, angulis anticis haud productis, deflexis, posticis rotundatis; elytris striatis, punctis seriatis grossis, interstitiis planis versus apicem multo angustioribus et convexioribus, 5° et 7° ante apicem conjunctis, 6° multum abbreviato; tarsis posticis elongatis.

Long. 3 lin. ♂ ♀.

Hab. PANAMA, Volcan de Chiriqui, David, Caldera (*Champion*).

10. *Selenophorus subsinuatus*.

Selenophorus subsinuatus, Putzeys, Stett. ent. Zeit. 1878, p. 31¹.

Hab. MEXICO, Cordova¹, Playa Vicente, Yolos (*Sallé*), Jalapa, Oaxaca (*Höge*); GUATEMALA, San Gerónimo, La Tinta, Dueñas, Rio Naranjo, Paraiso (*Champion*); NICARAGUA, Chinandega (*Sallé*); COSTA RICA, Cache (*Rogers*).

The form of the thorax varies greatly in the long series I have examined of this species; the hind angles are sometimes very obtuse, especially in a few of the Guatemalan examples; but the same form occurs in Oaxaca specimens, and it is impossible to separate them into species or local varieties.

11. *Selenophorus striatopunctatus*.

Selenophorus striatopunctatus, Putzeys, Stett. ent. Zeit. 1878, p. 33¹.

Hab. MEXICO, Chiapas¹, Cordova (*Sallé*), Oaxaca (*Höge*).—WEST INDIES¹.