

12. *Selenophorus flavipes*.

Selenophorus flavipes, Putzeys, Stett. ent. Zeit. 1878, p. 39¹.

Hab. MEXICO, Cordova (*Sallé*¹).

13. *Selenophorus semirufus*.

S. rufulo similis et affinis, sed multo major, sanguineus vel castaneo-rufus, abdomine, metathorace elytris (epipleuris solum exceptis) nigris; thorace convexo, medio quam elytra latiore, lateribus arcuatis, angulis anticis productis, posticis subrotundatis, basi ruguloso-punctulata; elytris apice vix sinuatis, apud suturam obtuse conjunctim rotundatis, dorso sericeo-nitentibus, argute subtiliter striatis, interstitiis planis 3^o-5^m et 7^o (juxta strias) conspicue pluripunctatis, omnibus versus apicem angustioribus striisque latoribus, striola scutellari elongata.

Long. 3 $\frac{3}{4}$ -4 $\frac{1}{4}$. ♂ ♀.

Hab. MEXICO, Guanajuato (*Dugès, coll. Sallé*).

14. *Selenophorus assimilis*.

Selenophorus assimilis, Putzeys, Stett. ent. Zeit. 1878 (January), p. 44¹.

Selenophorus mitis, Bates, Proc. Zool. Soc. 1878 (May), p. 591².

Hab. MEXICO, Oaxaca¹, Parada, Capulalpam (*Sallé*), Jalapa, Oaxaca (*Höge*), Mexico city (*Flohr*²); GUATEMALA (*Sallé*), near the city, San Gerónimo (*Champion*).

Distinguished by M. Putzeys by very slight and uncertain characters from the United-States *S. gagatinus*, Dej.

15. *Selenophorus tarsalis*. (Tab. III. fig. 23.)

Selenophorus tarsalis, Putzeys, Stett. ent. Zeit. 1878, p. 49¹.

Hab. MEXICO, Cordova¹, Jalapa (*Sallé*), Las Vigas, Oaxaca (*Höge*); BRITISH HONDURAS, river Sarstoon (*Blancaneaux*); GUATEMALA (*Sallé*), near the city, San Gerónimo, Balheu, San Isidro (*Champion*); COSTA RICA, Irazu (*Rogers*).

16. *Selenophorus splendidus*. (Tab. III. fig. 19.)

Selenophorus splendidus, Putzeys, Stett. ent. Zeit. 1878, p. 64¹.

Hab. MEXICO¹, Cordova, Yolos (*Sallé*); GUATEMALA¹.—SOUTH AMERICA, Colombia¹.

A specimen from Yolos, which I refer to this species, on account of the thorax narrowing towards the head, differs from the type in colours, being brassy green with opalescent lustre. It seems intermediate between *S. splendidus* and *S. tarsalis*.

17. *Selenophorus quadricollis*.

Selenophorus quadricollis, Putzeys, Stett. ent. Zeit. 1878, p. 55¹.

Hab. MEXICO¹; GUATEMALA, San Juan, Tamahu (*Champion*).

Varies in the depth of the elytral striæ; in individuals with deep striæ the punctures are scarcely visible.