

1. **Stenomorphus rufipes.** (Tab. III. fig. 22.)

*Stenomorphus rufipes*, Leconte, Proc. Ac. Phil. 1858, p. 59; id. Journ. Ac. Phil. iv. 1858, p. 29<sup>1</sup>, t. 4. f. 5.

*Hab.* NORTH AMERICA, Texas<sup>1</sup>.—MEXICO, Guanajuato (*Dugès, coll. Sallé*).

Named *S. rufipes* in the Sallé collection, and agreeing with Leconte's description and figure. The Mexican species, however, is different from a species collected in Texas by Belfrage, and distributed by him to his correspondents, which has the hind tibiæ less perceptibly dilated.

## GYNANDROPUS.

*Gynandropus*, Dejean, Sp. Gén. Col. v. p. 817; Putzeys, Stett. ent. Zeit. 1878, p. 289; Horn, Trans. Am. Ent. Soc. ix. (1881) p. 180.

A dozen species of this well-marked genus are known, all American, ranging from beyond the southern tropic to the temperate parts of North America, and including the West Indies.

1. **Gynandropus mexicanus.**

*Gynandropus mexicanus*, Putzeys, Stett. ent. Zeit. 1878, p. 291<sup>1</sup>.

*Hab.* MEXICO<sup>1</sup>, Cordova (*Sallé*), Las Vigas, Jalapa (*Höge*); GUATEMALA, San Gerónimo, Coatepeque, Senahu (*Champion*).

The species varies in Guatemala in the form and punctuation of the thorax; and I doubt if the following, founded on similar differences, is really distinct.

2. **Gynandropus intermedius.**

*Gynandropus intermedius*, Putzeys, Stett. ent. Zeit. 1878, p. 293<sup>1</sup>.

*Hab.* MEXICO, Oaxaca<sup>1</sup>, Guanajuato (*Dugès, coll. Sallé*).

3. **Gynandropus agonoides.**

*Gynandropus agonoides*, Putzeys, Stett. ent. Zeit. 1878, p. 294.

*Hab.* MEXICO.

Unknown to me.

## ATHROSTICTUS.

*Athrostictus*, Bates, Proc. Zool. Soc. 1878, p. 592.

*Hypolithus* (partim), Dejean.

*Selenophorus* (partim), Putzeys.

“Corpus elongato-ovatum, supra (præcipue elytris) dense minute punctatum. Caput ovatum, postice gradatim paullo angustatum. Oculi magni sed non prominentes, postice orbitu inclusi. Antennæ filiformes, graciles. Mentum profunde emarginatum, sinu fundo haud dentatum. Ligula angusta, apice libera, longe bisetosa; paraglossis latis ea longioribus. Elytra haud conspicue seriato-punctato. Cætera ut in generibus *Hypolitho* et *Harpalo*.”

BIOL. CENT.-AMER., Coleopt., Vol. I. Pt. 1, February 1882.

kk