

although the thorax is narrower and generally much more narrowed behind, I doubt whether it is specifically distinct.

4. *Bradycellus limbicollis*.

Oblongus, subdepressus, testaceo-fulvus vix nitidus, fronte, thoracis lateribus elytrorumque disco posteriore lateribusque nigro-fuscis; palpis, antennis basi pedibusque flavo-testaceis; thorace cordato-quadrato, postice modice rotundato-angustato, lateribus ante basin breviter sed fortiter sinuatis, angulis posticis productis, foveis basalibus latis plus minusve punctatis; elytris apice obtusis perparum sinuatis, margine basali utrinque fortiter arcuato, dorso argute striato, striola scutellari nulla, striis 8^a et 9^a ante basin approximatis, interstitiis planatis, 3^o unipunctato.

Long. $1\frac{3}{4}$ lin.

Hab. MEXICO (*Sallé*); GUATEMALA, Paso Antonio (*Champion*).

Var. omnino testaceo-flava (*immaturus*?).

Hab. GUATEMALA, Paso Antonio (*Champion*).

5. *Bradycellus nigrellus*.

B. circumdato colore similis, multo minor, piceo-niger (elytris interdum subæneo tinctis), nitidus, collo interdum rufo; palpis, labro, antennis basi, pedibusque flavo-testaceis, elytris margine suturaque (præcipue postice) rufo-testaceis; thorace quadrato postice recte modice angustato, antice parum rotundato, angulis posticis rectis, basi transversim depressa, fovea utrinque lata discrete fortiter punctata; elytris subparallelis, apice obtuse rotundatis, striatis, striola scutellari nulla, interstitiis planatis, 3^o unipunctato.

Long. $1\frac{1}{2}$ lin.

Hab. MEXICO, Cordova (*Sallé*); GUATEMALA, near the city, Quiche Mountains, San Gerónimo, Capetillo, Zapote, Panajachel (*Champion*).

6. *Bradycellus angulifer*.

B. nigrello quam maxime affinis, differt colore fulvo-castaneo, elytris utrinque disco, thoraceque dorso interdum, infuscatis. Parvus; thorace subcordato, antice sat rotundato-dilatato, postice magis angustato, margine ante angulos productos acutos conspicue sed breviter sinuato, basi utrinque punctata; elytris absque striola scutellari; antennis basi, palpis pedibusque testaceo-flavis.

Long. $1\frac{1}{3}$ lin.

Hab. MEXICO, Vera Cruz, Cordova (*Sallé*), Guanajuato (*Dugès, coll. Sallé*), Oaxaca (*Höge*); GUATEMALA, Capetillo, San Gerónimo (*Champion*).

I adopt the name given to this species by Chaudoir in the *Sallé* collection. It is scarcely distinct from *B. nigrellus*, differing only in colour and in the thorax being a little more rounded anteriorly and more cordate. As all the Mexican specimens I have seen are of this form, to the exclusion of the black *B. nigrellus*, it merits a distinguishing name.

The following genus, in the present state of our knowledge, is difficult to place in the Harpalidæ series. It has hitherto been associated with *Acinopus*, *Daptus*, and other genera in a distinct subgroup; but, as Dr. Horn points out, the penultimate joint