

attingente) usque ad suturam paullo ante apicem ducta, relictā macula magna communi triangulari suturali cum margine basali per vittam suturalem connexa, et limbo toto, fusco-nigris, margine ipso laterali et apicali fulvo; thorace sicut in *P. bifasciata* valde transverso, margine ab angulo mediano usque ad basin paullulum flexuoso; elytris amplis, subtilissime punctulato-striatis, interstitiis convexis juxta apicem altioribus; pectore et abdomine fusco-nigris.

Long.  $2\frac{3}{4}$  lin.

*Hab.* PANAMA, Volcan de Chiriqui (*Champion*).

One example only.

#### 4. *Pentagonica omostigma*. (*Pentagonica homostigma*, Tab. IX. fig. 23.)

*P. gonostigmæ* affinis et similis, sed elytrorum signaturis aliis etc. Testaceo-fulva, antennarum articulis 4 basalibus fusco-maculatis elytrisque utrinque macula rotundata humerali et fascia lata ante apicem (ad suturam et marginem dilatata) nigris, margine omnino testaceo-fulvo; capite subopaco; thorace valde transverso, margine ab angulo mediano usque ad basin flexuoso; elytris distinctius exarato-striatis, striis punctatis, interstitiis sat convexis, 9° (marginali) catenato-punctato.

Long.  $2\frac{1}{2}$  lin.

*Hab.* PANAMA, Volcan de Chiriqui, Bugaba (*Champion*).

#### 5. *Pentagonica picticornis*. (Tab. IX. fig. 25.)

*P. gonostigmæ* iterum affinis. Parum convexa; capite supra et infra nigro; antennis nigris, articulis 4 basalibus rufis, 2°–4<sup>m</sup> pallidioribus; thorace rufo valde transverso, postice minus rapide angustato margineque valde flexuoso; elytris elongato-ovatis, striis punctulatis vix impressis sed interstitiis convexis, 9° (marginali) catenato-punctato, fusco-nigris, vitta lata humerali margineque toto testaceo-rufis; abdomine postice fusco-nigro.

Long.  $2\frac{1}{2}$  lin.

*Hab.* GUATEMALA, El Jicaró (*Champion*).

One example only.

#### 6. *Pentagonica semifulva*.

Rufescenti-fulva, elytris (margine excepto), meso- et metathorace abdomineque nigris, palpis et antennis nigro-fuscis; thorace transverso postice minus rapide angustato, margine mox pone angulum prominentem recto vel subsinuato, limbo antico sæpe infuscato; elytris vix impresso-striatis, sed interstitiis sat convexis, 9° catenato-punctato, femoribus et tibiis anticis sæpe subinfuscatis.

Long. 2– $2\frac{1}{3}$  lin.

*Hab.* MEXICO, Cordova (*Sallé*), Jalapa (*Höge*); GUATEMALA, Cerro Zunil, Pantaleon (*Champion*).

This must approach very closely *P. bicolor*, Leconte, of the Western United States, and may be the same species; but the description does not mention that the meso- and metathorax are black as well as the abdomen.

#### 7. *Pentagonica maculicornis*.

*P. semifulvæ* simillima, differt tantum antennis fulvis, scapo toto cæterisque articulis apice solum fuscis;

BIOL. CENTR.-AMER., Coleopt., Vol. I. Pt. 1, September 1883.

2 ff