

7. *Loxopeza cyane*. (Tab. X. fig. 3.)

L. majusculæ (Chd.) (*L. grandis*, Hald.?) quoad formam et sculpturam simillima, sed differt elytris saturatius cæruleis subviolaceis, minus nitidis. Testaceo-rufa, antennarum articulis 4^o–11^m fuscis, ventre (processu intercoxali excepto) nigro, elytris saturate cæruleis minus nitidis; capite fere lævi; thorace transverso, antice sat late rotundato, postice subrecte, modice sed distincte angustato, angulis posticis perparum productis, disco subtiliter vage striguloso; elytris quam in *L. majuscula* longius oblongis, acute striatis, interstitiis modice convexis, versus latera deplanatis; tarsorum posticorum articulo penultimo emarginato, angulis vix elongatis.

Long. $4\frac{1}{2}$ lin.

Hab. MEXICO, Guanajuato (*Dugès*, coll. *Sallé*), Mexico city (*Flohr*, *Höge*).

Closely allied to the Texan *L. majuscula* (Chaud.), which, I understand Dr. Horn, refers to the prior-described *L. grandis* of Haldeman. It is, however, sufficiently distinct by its rather more elongate form and the different colour of its elytra, the Texan species being bright brassy green with a bluish tinge. I have compared together half a dozen examples of each.

8. *Loxopeza rufosutura*.

Loxopeza rufosutura, Chaudoir, Monogr. des Lébiides, p. 31¹.

Lebia rufosutura, Motschulsky, Bull. Mosc. 1864, ii. p. 225².

Hab. MEXICO (*Sallé*¹), Vera Cruz, Cordova (*Sallé*), Oaxaca, Juquila (*Boucard*, coll. *Sallé*), Jalapa (*Höge*); GUATEMALA¹; NICARAGUA².

I adopt the name as given to specimens in the *Sallé* collection; but the description of Motschulsky seems scarcely to apply to the Mexican examples (the only ones I have seen), not one of which has a red suture.

9. *Loxopeza chloroptera*.

Loxopeza chloroptera, Chaudoir, Monogr. des Lébiides, p. 32¹.

Lebia chloroptera, Chaudoir, Ann. Soc. Ent. Fr. 1835, p. 437.

Lebia nigriventris, Putzeys, Mém. Soc. R. des Sci. de Liège, 1846, ii. p. 382².

Hab. NORTH AMERICA, FLORIDA? (*Reiche*¹).—MEXICO^{1 2}, Ventanas (*Forrer*), Guajuco in Nuevo Leon (*Dr. Palmer*), Guanajuato (*Dugès*, coll. *Sallé*), Mexico city (*Flohr*), Tlacotalpam, Oaxaca (*Boucard*, coll. *Sallé*), Yucatan (*Pilate*¹).

According to the series in the *Sallé* collection and the specimens collected by others in different parts of Mexico, this appears to be a very variable species, at least as far as regards the colour of the elytra and the more or less reddish hue of the epipleuræ and margins. The rather broader thorax seems to be the only character which separates it from *L. rufosutura*.

10. *Loxopeza atriventris*.

Lebia atriventris, Say, Trans. Amer. Phil. Soc. ii. p. 13¹ (1823); Dejean, Sp. Gén. Col. ii. p. 454.

Loxopeza atriventris, Chaudoir, Monogr. des Lébiides, p. 32².

Hab. NORTH AMERICA, Eastern States¹.—MEXICO².