

11. **Loxopeza melanocephala.**

Loxopeza melanocephala, Chaudoir, Monogr. des Lébiides, p. 33¹.

Hab. MEXICO¹, Cuernavaca, Yolos, Durasnal, Oaxaca (*Boucard, coll. Sallé*), Mirador, Jalapa (*Höge*).

12. **Loxopeza rufolimbata.**

Loxopeza rufolimbata, Chaudoir, Monogr. des Lébiides, p. 33¹.

Hab. MEXICO, Cuernavaca (*Boucard, coll. Sallé*¹), Guanajuato (*Dugès, coll. Sallé*), Cordova (*Sallé*), Jalapa (*Höge*).

13. **Loxopeza exarata.** (Tab. X. fig. 5.)

L. melanocephalæ quoad colores similis, sed differt elytrorum striis usque ad apicem et latera profunde insculptis interstitiis æqualiter et valde convexis etc. Oblongo-ovata, testaceo-rufa, capite supra, genis lateribus abdomineque nigris, elytris obscurius viridi-æneis, antennarum articulis 4^o–11^m interdum piceis; capite longitudinaliter strigoso, vertice sublævi; thorace fere semiovato, antice rotundato (haud ampliato), postice haud vel perparum angustato, angulis posticis rectis, disco subtiliter strigoso; epipleuris versus basin rufo-piceis; tarsorum posticorum articulo penultimo emarginato.

Long. 3½ lin.

Hab. MEXICO, Jalapa (*Höge*).

Var. yoloensis. Caput supra grossius strigosum et punctatum, infra nigrum linea angusta mediana rufa; thorax crebre coriaceo-strigosus, minus nitidus, angulis posticis paullo exstantibus.

Hab. MEXICO, Yolos (*Boucard, coll. Sallé*).

14. **Loxopeza eburata.** (Tab. X. fig. 6.)

L. obliquatæ affinis. Breviter oblongo-ovata, testaceo-rufa, nitida, antennarum articulis 3^o–11^m femorum apice, tibiis et tarsis (unguiculis rufis exceptis) elytris nigris, his utrinque vitta eburnea, a basi interstitii 5ⁱ usque fere ad apicem interstitii 4ⁱ ducta; palpis piceis, apice rufis; capite fere lævi; thorace valde transverso (duplo magis lato quam longo), antice late rotundato fere angulato, postice haud angustato lateribus et angulis posticis rectis, margine laterali explanato-reflexo latissimo, dorso vage ruguloso; elytris profunde striatis, interstitiis valde convexis, quarto quam cætera latiore; tarsorum posticorum articulo penultimo emarginato.

Long. 3¼ lin. ♀.

Hab. PANAMA, Volcan de Chiriqui (*Champion*).

The white stripe on each elytron runs for a short distance from the base along the fifth interstice; it then leaves the fifth for the fourth, on which it commences lower down than it did on the fifth, and continues very nearly to the apex.

METABOLA.

Metabola, Chaudoir, Monogr. des Lébiides, p. 50.

The following species is the only one at present known of this genus.

1. **Metabola rufopyga.**

Metabola rufopyga, Chaudoir, Monogr. des Lébiides, p. 50¹.

Hab. MEXICO¹, Cordova, Capulalpam, Coscomatepec (*Sallé*), Mirador (*Höge*).