

Chaudoir's description as far as colour is concerned. There is, however, some ground for considering it an extreme variety of *C. viridis*, as the blue form of the species in Mexico is smaller and more brilliant than the northern type, and is subject to variations in colour.

10. *Lebia dugesi*.

L. viridi simillima, sed differt antennarum articulis 1^o-3^m testaceis, thoracisque angulis posticis productis dentiformibus. Oblonga, cærulea, elytris profundius punctulato-striatis, interstitiis subconvexis.

Long. 5 millim.

Hab. MEXICO, Guanajuato (*Dugès, coll. Sallé*).

One example only.

11. *Lebia ignita*. (Tab. X. fig. 10.)

L. viridi affinis, sed relative brevior, thorace præcipue valde transverso colloque supra depresso. Læte cærulea, elytrorum dimidio apicali rufo-aureo splendidissimo.

Long. 4-4½ millim.

Hab. GUATEMALA, Dueñas, Capetillo, Cerro Zunil (*Champion*).

Distinctly shorter and relatively broader than *L. viridis*, the thorax especially being at least twice as broad as long (excluding the basal lobe), and the margin near the hind angles more widely explanated. The crown of the head is also distinctly convex, or the neck depressed. The elytra are similarly striated, with flat interstices. The colour is a rich blue, with the apical half of the elytra fiery golden purple. Tarsi bilobed as in *L. viridis*.

12. *Lebia cœlina*. (Tab. X. fig. 11.)

Parva, late ovata, convexa, saturate cærulea (elytris interdum violaceis), antennis articulis basalibus labroque piceis, palpis acutis; capite brevi, lævi, post oculos cito angustato sed collo supra haud depresso; thorace transverso (latiore quam in *L. viridi*), trapezoidali, postice usque ad angulos valde acutos paullo dilatato, lateribus subrectis, margine basali prope angulum sinuato; elytris ampliatis subtilissime striatis; tarsorum articulo penultimo elongato-bilobato.

Long. 3 millim.

Hab. PANAMA, Volcan de Chiriqui (*Champion*).

Belongs to the *L. viridis* group, but distinguished by its short plump form and the trapeze-like outline of the short and broad thorax.

13. *Lebia xanthogona*.

L. pumilæ similis, sed differt thorace multo latiore angulisque posticis fulvo-testaceis etc. Breviter ovata, ænescenti-fusca nitida, antennarum articulis 3 basalibus, pedibus (femoribus medio tarsisque infuscatis) thoracisque angulis posticis pallide fulvo-testaceis; capite lævi; thorace valde transverso, postice paullo angustato (lateribus subrectis), angulis posticis vix rectis, margine basali juxta angulum obliquo recto; elytris subtiliter punctulato-striatis, interstitiis subconvexis; tarsorum articulo penultimo bilobato.

Long. 4 millim.

Hab. MEXICO, Orizaba (*Sallé*), Jalapa (*Höge*).