

14. **Lebia agnata.**

Lebia agnata, Chaudoir, Monogr. des Lébiides, p. 86¹.

Hab. MEXICO¹, Orizaba, Puebla (*Sallé*), Mirador, Jalapa (*Höge*).

15. **Lebia croceicollis.** (Tab. X. fig. 13.)

L. pumilæ affinis, quoad formam *L. xanthogonæ* simillima, sed differt colore nigro (capite elytrisque subænescentibus), palpis nigro-fuscis, antennarum articulis $3\frac{1}{2}$ basalibus thoraceque croceis, pedibus piceo-fuscis, femoribus 4 anticis basi pallidis; capite ovato, lævissimo, post oculos gradatim rotundato-angustato; thorace valde transverso, postice paullulum angustato, angulis posticis paullo exstantibus, margine laterali angusto subrecto; elytris obsolete punctulato-striatis; tarsis bilobatis.

Long. $3\frac{1}{2}$ millim.

Hab. MEXICO, Jalapa (*Höge*).

Var. Femora et tibiæ pallide testacea, elytrorumque interstitia paullulum convexa.

Long. $4\frac{1}{4}$ millim.

Hab. GUATEMALA, San Gerónimo (*Champion*).

One example only.

16. **Lebia goniessa.** (Tab. X. fig. 12.)

Parva, breviter ovata, convexa, nigra, nitida; capite, thorace, antennarum articulis 4 basalibus (cæteris fuscis) pedibusque fulvo-rufis, femoribus posticis dimidio basali tibiisque dimidio apicali sæpissime nigro-fuscis; capite ovato lævissimo; thorace vix transverso trapezoideo (postice dilatato), margine recto antice angusto postice valde dilatato angulisque retrorsum productis, margine basali prope angulos paullo retrorsum obliquo; elytris subtilissime punctulato-striatis; tarsis bilobatis.

Long. $3\frac{3}{4}$ –4 millim.

Hab. GUATEMALA, San Gerónimo (*Champion*).

Belongs to the *viridis* group, and allied to *L. agnata*; differing from other species with red head and thorax in the shape of the hind thoracic angles, which are produced backward almost as in *L. cœlina*. The head is perfectly free from striæ; and the striæ of the elytra are scarcely impressed, although the interstices are slightly convex. The elytra with margins and epipleuræ and the abdomen are deep black. The two punctures of the third interstice are small, but very distinct, and situated in the third stria.

17. **Lebia microtes.**

L. goniessæ quam maxime similis et affinis, sed thoracis angulis posticis rectis nec retrorsum productis, fronteque striato; Parva, nigra, nitida, capite, antennis, thorace et pedibus rufo-fulvis, palpis nigris; capite ovato, fronte præcipue prope oculos subtiliter pluristrigoso; thorace valde transverso postice, haud dilatato, angulis posticis rectis ibique margine explanato-reflexo, margine basali prope angulos recto; elytris breviter ovatis, striis distinctius et acutius impressis.

Long. 3 millim.

Hab. GUATEMALA, Panima, Chacoj, Aceituno, Volcan de Atitlan (*Champion*).

It is difficult to believe this is more than a local modification of *L. goniessa*; the differences of form and sculpture, however, indicated in the above diagnosis appear to