

formi humerali, et 1 postmediana transversa, antice apud suturam indentata, latera haud attingente, apice autem anguste nigro.

“Long. $3\frac{1}{4}$ lin.”

Hab. MEXICO, Cordova (*Sallé*); GUATEMALA, Pantaleon (*Champion*); PANAMA, Volcan de Chiriqui, David, Bugaba (*Champion*); NICARAGUA, Chontales (*Belt*¹).

28. *Lebia clio*. (Tab. X. fig. 19.)

L. variegata (Dej.) quoad colores et signaturam simillima, sed macula elytrorum suturali-apicali deficiente. Flavo-testacea (abdomine apice concolore), capite (lævi) supra et infra, palpis antennisque nigris his articulis 1^o–3^m et 4^o basi flavis; thorace transverso, postice parum angustato, angulis posticis sat productis, margine basali juxta angulum rotundato-obliquo; elytris subtiliter punctato-striatis, interstitiis passim planis, flavis, macula circumscutellari, altera subvirguliformi humerali, vitta suturali fasciaque lata undulata versus apicem nigris, sutura post fasciam haud nigra.

Long. $6\frac{1}{2}$ millim.

Hab. MEXICO, Playa Vicente (*Sallé*), Jalapa (*Höge*).

Extremely near *L. variegata* (Dej.), but the striæ much less impressed, the interstices flat, and the apical sutural black spot wanting.

29. *Lebia variegata*.

Lebia variegata, Dej. Sp. Gén. Col. v. p. 375; Chaudoir, Monogr. des Lébiides, p. 91¹.

Hab. MEXICO, Jalapa (*Höge*); GUATEMALA, Las Mercedes (*Champion*); PANAMA, Bugaba, David, Caldera (*Champion*).—SOUTH AMERICA, Colombia¹.

In all the specimens from Jalapa the black sutural stripe terminates at the posterior fascia, and the apical sutural spot is reduced to a narrow black apical edging; the palpi also are darker. This variety forms a transition, therefore, to the *L. clio* of the same locality. The concave pygidium is sometimes blackish both in the Mexican and Panama forms.

30. *Lebia interrupta*.

Lebia interrupta, Chaudoir, Monogr. des Lébiides, p. 100¹.

Hab. MEXICO¹.

I have not seen this species, which M. de Chaudoir obtained from M. E. Deyrolle.

31. *Lebia anchora*. (Tab. X. fig. 20.)

Lebia anchora, Chevrolat, Col. Mex. ii. no. 132 (1835); Chaudoir, Monogr. des Lébiides, p. 102¹.

Lebia bonellii, Putzeys, Mém. Soc. R. de Liège, ii. p. 391 (1846).

Hab. MEXICO¹, Vera Cruz, Tuxtla, Yolos (*Sallé*), Mirador (*Höge*); GUATEMALA, Capetillo, San Gerónimo (*Champion*).