

32. *Lebia incommoda*.

Lebia incommoda, Chaudoir, Monogr. des Lébiides, p. 103¹.

Hab. MEXICO, Campeche (*Pilate*¹).

33. *Lebia scalpta*. (Tab. X. fig. 21.)

L. subrugosa (Chaud.) similis, sed differt capite a collo usque ad epistoma subrecte multistriato. Oblonga, flavo-testacea, capite supra et infra elytrisque signaturis nigris; labro, partibus oris antennisque rufo-testaceis; capite supra acute striato, striis fere omnibus rectis a collo usque ad epistoma; oculis magnis prominentibus; thorace sat transverso, marginibus late explanato-reflexis; elytris oblongis, profunde striatis, interstitiis convexis, fascia ante apicem triloba vel lacerata, variabili, vitta submarginali a fascia usque ad humerum gradatim attenuata, macula quadrata circumscutellari vittaque suturali inter maculam scutellarem et fasciam nigris; tarsorum 2 posticorum articulo 4^o profunde emarginato, 4 anteriorum bilobatis.

Long. 6½ millim.

Hab. MEXICO, Jalapa (*Höge*), Yucatan (*Sallé*).

Similar in the black markings of the elytra to *L. subrugosa*, Chaud., but differing in the uniform nearly straight sculpture of the head. The posterior black fascia is, in most examples, trilobed almost as regular as in *L. variegata* and *L. anchora*, but in others it is irregular and ragged on both its edges—a specimen, wrongly named *L. subrugosa*, in the Sallé collection reaching an extreme in this respect, and agreeing, therefore, with Chaudoir's figure of *L. subrugosa*.

34. *Lebia duillia*. (Tab. XI. fig. 1.)

L. striaticipiti (Chaud.) affinis. Oblongo-ovata, postice dilatata, flavo-testacea (capite thoraceque rufioribus), elytris subcyaneo-nigris utrinque maculis duabus magnis marginibusque angustis flavo-testaceis, altera prope basin oblique quadrata a stria 1^a usque ad striam 7^m extensa, interdum usque ad marginem extensa, altera apicali transversa (sutura apiceque suturali fuscis); capite supra longitudinaliter striato, spatio triangulari prope epistoma lævi; thorace brevi et lato, marginibus late explanatis, ante angulos leviter sinuatis, dorso subtiliter (præcipue longitudinaliter) strigoso; elytris profunde striatis, interstitiis convexis prope apicem deplanatis; tarsorum 4 anter. articulo 4^o bilobato, 2 poster. lobis brevioribus.

Long. 4½ millim.

Hab. MEXICO, Jalapa (*Höge*).

The striated and red-coloured head brings this species into the same subsection as *L. ægra*, *striaticeps*, &c., but in the elytral markings it very much resembles *L. chiri-quensis*, *nemoralis*, and allies; it wants, however, the long linear lobes of the fourth joint of the hindmost tarsi, which distinguish the latter subgroup. The colour of the elytral markings is deep black, sometimes with a bluish tinge.

35. *Lebia subrugosa*.

Lebia subrugosa, Chaudoir, Monogr. des Lébiides, p. 118, t. 1. fig. 49¹.

Hab. MEXICO¹, Capulalpam (*Sallé*), Jalapa (*Höge*); GUATEMALA, San Gerónimo (*Champion*).