

40. **Lebia soror.** (Tab. XI. fig. 2.)

Lebia soror, Chaudoir, Monogr. des Lébiides, p. 122, t. 1. fig. 54¹.

Hab. MEXICO, Cordova (*Sallé*); NICARAGUA, Chontales (*Belt*); PANAMA, Las Mercedes, Caldera, Bugaba (*Champion*).—SOUTH AMERICA, Amazons¹.

41. **Lebia centromaculata.** (Tab. XI. fig. 3.)

Lebia centromaculata, Putzeys, Mém. Soc. R. d. Sci. de Liège, ii. p. 386; Chaudoir, Monogr. des Lébiides, p. 124, t. 2. fig. 59¹.

Hab. MEXICO, Juquila (*Sallé*, *Höge*); GUATEMALA (*Sallé*), Capetillo, San Gerónimo (*Champion*).

Var. *L. miradora*. Elytris sutura nigra a scutello usque ad maculam suturalem fasciaque mediana usque ad interstitium 9^m extensa, apud interstitia 6^m–9^m dilatata et retrorsum inflexa.

Hab. MEXICO, Mirador (*Höge*).

42. **Lebia arietis.** (Tab. XI. fig. 4.)

L. centromaculatæ proxime affinis et similis; thorace ejusdem formæ, sed elytris fascia tenui valde angulata (apud suturam haud maculam efficiente) suturaque antice nigra usque ad scutellum.

Long. 5½ millim.

Hab. PANAMA, Volcan de Chiriqui (*Champion*).

The thorax is of the same form as in *L. centromaculata*, *i. e.* much rounded from the anterior angles, and thence slightly narrowed in nearly a straight line to the base; but the black mark of the elytra in all the very numerous examples differs constantly in not consisting of a rhomboidal spot on the suture with a slender angular branch on each side, but forming an angular line or fascia of equal width throughout, the pale ground-colour at the suture behind indenting nearly always the black mark. The elytra also seem more deeply striated, the striæ alveolated and darker than the interstices.

43. **Lebia chelostigma.** (Tab. XI. fig. 5.)

Fulvo-testacea, elytris macula communi pone medium utrinque ramum rectum antrorsum obliquum emittente, suturaque fere usque ad scutellum, nigris; capite antice, cum mandibulis et maxillis elongato, collo angusto, oculis valde prominentibus; antennis elongatis, gracilibus; thorace antice (a basi) curvatim angustato, marginibus (præcipue postice) explanato-reflexis, angulis posticis rectis, dorso subtiliter alutaceo; elytris elongato-ovatis sat convexis, profunde striatis, interstitiis convexis; tarsorum articulo 4^o bilobato.

Long. 6–8 millim.

Hab. PANAMA, Volcan de Chiriqui, Bugaba (*Champion*).

In the markings of the elytra similar to *L. centromaculata*, but differing from that species much in the form of the thorax, the widest part of which is at the hind angles, and the sides not being rounded outward immediately behind the anterior angles. The prolongation of the anterior part of the head with the mandibles and maxillæ is visible chiefly in the larger examples and in the males.