

### 70. *Lebia intermedia*.

*Dianchomena intermedia*, Chaudoir, Monogr. d. Lébiides, p. 199, t. 3. fig. 142<sup>1</sup>.

*Hab.* MEXICO, Cordova (*Sallé*), Yucatan (*Pilate*<sup>1</sup>).

### 71. *Lebia calliparis*. (Tab. XII. fig. 1.)

*L. intermediae* affinis, differt elytris bifasciatis. Parva, flavo-testacea, capite, palpis, antennis (articulis 1<sup>o</sup>–3<sup>m</sup> exceptis), fasciis elytrorum duabus, prima lata basali, postice fere recta, secunda lata subtrilobata suturaque angusta, nigris; capite postice valde convexo collo constricto, fronte media lævi, lateribus striatis; thorace rufo-testaceo, transverso, antice rotundato, lateribus explanato-reflexis; elytris obsolete striatis, suturali solum acute impressis, interstitiis planis; tarsorum omnium articulo quarto anguste bilobato.

Long. 4½ millim.

*Hab.* GUATEMALA, Capetillo (*Champion*).

### 72. *Lebia abdominalis*.

*Lebia abdominalis*, Chaudoir, Bull. Mosc. 1843, p. 704; Leconte, Ann. Lyceum N. York, iv. p. 195 (1848)<sup>1</sup>.

*Dianchomena abdominalis*, Chaud. Monogr. d. Lébiides, p. 191<sup>2</sup>.

*Hab.* NORTH AMERICA, Alabama<sup>1</sup>, Louisiana<sup>2</sup>.—MEXICO<sup>2</sup>, Guanajuato (*Dugès*, coll. *Sallé*); BRITISH HONDURAS (*Blancaneaux*); GUATEMALA, Champerico (*Champion*); NICARAGUA, Chontales (*Belt*).

This with two South-American species are the typical forms of Chaudoir's genus *Dianchomena*; but *L. abdominalis* is obviously nearly related to the *L. quadricolor* section of true *Lebiæ*.

### 73. *Lebia vittata*.

*Carabus vittatus*, Fabr. Syst. El. i. p. 202.

*Lebia vittata*, Dej. Sp. Gén. Col. i. p. 267<sup>1</sup>.

*Lebia flavovittata*, Chevrolat, Col. Mex. ii. no. 161<sup>2</sup>.

*Aphelogenia vittata*, Chaudoir, Monogr. d. Lébiides, p. 184, t. 3. fig. 127<sup>3</sup>.

*Hab.* NORTH AMERICA, United States, generally distributed<sup>1 3</sup>.—MEXICO<sup>2 3</sup> (*Sallé*).

This and the following two species, forming part of Chaudoir's Section II. of *Aphelogenia*, form a tolerably natural group very distinct from his Section I.

### 74. *Lebia histrionica*. (Tab. XII. fig. 2; fig. 3, var. *c*.)

*L. vittatæ* quoad formam similis, sed striis impressis interstitiisque paullo convexis. Flavo-testacea, capite thoraceque rufis, palpis, antennis (basi exceptis), femoribus (basi excepta), genibus, tibiis apice et tarsis, nigris; elytris signaturis nigris variabilibus; capite punctulato, fronte utrinque strigosa; thorace quam in *L. vittata* paullo latius explanato-marginato, disco transversim strigoso; tarsorum posticorum articulo quarto breviter bilobato.

Long. 6 millim.

*Hab.* MEXICO (*Sallé*); GUATEMALA (*Champion*).