

Var. *a* (*L. scutellata*, Chevr. MS.). Elytris macula quadrata (postice haud angulata) circumscutellari, fasciaque postmediana apud suturam latissima versus latera angustata, nigris.

Hab. MEXICO, Playa Vicente (*Sallé*).

Var. *b*. Macula circumscutellari propter suturam flavescentem fere divisa, fascia posteriore utrinque in maculas duas divisa, interstitiis tertio quartoque flavis.

Hab. MEXICO, Cordova (*Sallé*).

Var. *c* (*L. nigrosignata*, Dugès). A var. *b* differt solum parte mediana fasciæ posterioris usque ad maculam circumscutellarem continuata.

Hab. MEXICO, Guanajuato (*Dugès, coll. Sallé*).

Var. *d*. Macula circumscutellari deficiente, fascia posteriore utrinque divisa sed antice prolongata, parte suturali sicut in var. *c* vittiformi, parte discoidali in maculam antice acuminatam extensa.

Hab. GUATEMALA, Toco (Champion).

75. *Lebia bivittata*.

Carabus bivittatus, Fabr. Ent. Syst. Suppl. p. 59.

Lebia quadrivittata, Dej. Sp. Gén., Col. i. p. 268; Leconte, Ann. Lyc. N. Y. iv. p. 195¹.

Aphelogenia bivittata, Chaudoir, Monogr. d. Lébiides, p. 189, t. 3. fig. 132.

Hab. NORTH AMERICA, Alabama¹, Texas.—MEXICO, Guanajuato (*Dugès, coll. Sallé*), near the capital (*Flohr*).

76. *Lebia quadriplagiata*.

Aphelogenia quadriplagiata, Chaudoir, Monogr. d. Lébiides, p. 172, t. 3. fig. 113¹.

Hab. MEXICO, Yucatan (*Pilate*¹).

77. *Lebia amabilis*.

Aphelogenia amabilis, Chaudoir, Monogr. d. Lébiides, p. 174, t. 3. fig. 117¹.

Hab. GUATEMALA, Tamahu, Cahabon (*Champion*); PANAMA, Volcan de Chiriqui (*Champion*).—SOUTH AMERICA, Colombia¹.

78. *Lebia hilaris*. (Tab. XII. fig. 6.)

Aphelogenia hilaris, Chaudoir, Monogr. d. Lébiides, p. 177, t. 3. fig. 120¹.

Hab. MEXICO, Cordova¹, Toxpam (*Sallé*).

79. *Lebia extrema*.

L. hilari et *amabili* affinis; differt thorace magis quadrato, postice haud dilatato, lateribus rectis, angulis posticis minus productis. Breviter ovata, convexa (minus quam *L. amabilis*), flavo-testacea; fasciis duabus elytrorum, prima basali (postice paullulum flexuosa), secunda mediana (antice paullo flexuosa), per vittam suturalem connexis, nigris; capite lævi, vertice palpisque infuscatis; thorace transversim quadrato, lateribus fere rectis, lobo basali maxime elongato; elytris modice convexis, apice leviter sinuatim truncatis, striis punctulatis vix impressis, interstitiis planis; pectore ventrisque basi fuscis.

Long. 3 millim.

Hab. PANAMA, Volcan de Chiriqui (*Champion*).