

80. *Lebia granaria*.

Lebia granaria, Putzeys, Mém. Ac. Sci. de Liège, ii. 392¹.

Aphelogenia granaria, Chaudoir, Monogr. de Lébiides, p. 174, t. 3. fig. 116.

Hab. PANAMA, Volcan de Chiriqui (*Champion*); SOUTH AMERICA, Venezuela¹.

An example from Panama resembles closely a specimen from Caraccas, sent me by M. René Oberthür.

The four preceding species form, with others from various parts of Tropical America, a natural group, one of the chief characteristics of which is the convex elytra.

81. *Lebia heraldica*.

Oblonga, flavo-testacea, elytris nigris, vitta utrinque flava a basi (imprimis per interstitium quintum, deinde interstit. 4^m et 3^m) usque ad fasciam subapicalem flavam extensa, marginibus angustis flavis, antennis articulis 5^o–11^m (et 4^o apice) fuscis; capite fere lævi; thorace late transverso, antice rotundato-dilatato, postice sat angustato, lateribus late explanato-reflexis, lobo basali subelongato quadrato, dorso lævi; elytris acute striatis, interstitiis paullo convexis apice planioribus; tarsorum omnium articulo quarto longe bilobato.

Long. 7 millim.

Hab. GUATEMALA, Cerro Zunil (*Champion*).

One example only, slightly immature.

Does not fit well in any of the subsections of Chaudoir's Monograph; but it is doubtless allied to the species of the *L. variegata* group.

82. *Lebia coptoderina*. (Tab. XII. fig. 4.)

Coptoderæ tripartitæ (Chaud.) subsimilis. Oblonga, parum convexa, flavo-testacea; elytris nigro-æneis fasciis duabus (prima valde dentata) flavis; capite lævi; thorace transverso, antice sat late rotundato, postice angustato, lateribus subæqualiter explanato-reflexis, dorso lævi; elytris postice paullo ampliatis, apice sinuatis, dorso profunde striatis, interstitiis convexis, fascia flava prima paullo ante medium, apud interstitium quartum postice, apud interstitium quintum antice, dilatata, fascia secunda juxta apicem antice undulata, postice ab apice margine viridi prope suturam separata; tarsorum posticorum articulo quarto breviter bilobato, lobis acuminatis; palpis tenuibus.

Long. 6 millim.

Hab. PANAMA, Volcan de Chiriqui (*Champion*).

One example only. The position of the species in Chaudoir's classification is uncertain.

83. *Lebia mesostigma*. (Tab. XII. fig. 5.)

Convexa, rufo-testacea, elytris macula parva suturali paullo post medium alteraque minore utrinque paullo posteriore versus marginem, nigris; capite sparsim subtiliter punctulato, prope oculos strigoso, collo sat lato et convexo usque ad oculos recto; palpis apice obtusis; thorace subrotundato basi perparum lobato, lateribus fere regulariter arcuatis, antice magis quam postice angustato, margine explanato-reflexo, tota superficie strigoso-coriacea; elytris vix truncatis fere obtuse rotundatis, profunde striatis, interstitiis valde convexis; tarsorum posticorum articulo quarto bilobato, lobis latis.

Long. 8 millim.

Hab. MEXICO, Tepanistlahuaca (*Sallé*).

One example only.