

colour of the tibiæ and antennæ. It is nearly allied also to *A. dominula* (Bates) from Peru, differing in the elytral striæ being scarcely impressed, and in colour.

2. *Agra insidiosa*.

A. eurypelmæ affinis; differt solum statura graciliori, antennis palpisque rufo-piceis, pedibus rufis. Ventris (♂) segmentis ultimo et penultimo medio longitudinaliter aciculatis, segmento antepenultimo medio crebre punctulato.

Long. 20 millim. ♂.

Hab. BRITISH HONDURAS (*Blancaneaux*).

One example only.

Agrees with *A. eurypelma* in every respect except its narrower form, red legs (including coxæ), dull red antennæ and palpi, and the different sculpture of the ventral segments of the male. The colour is rather more metallic.

3. *Agra castaneipes*. (Tab. XII. fig. 12.)

A. gemmatæ (Klug) valde affinis, quoad formam simillima, sed differt colore æneo (subtus subviridi-æneo) elytrorumque foveis fundo cyaneis; partibus oris, antennis pedibusque castaneo-rufis. Gracilis, ænea, polita; capite postice elongato apud collum solum angustato; thorace angusto, lineatim foveolato; elytris lineatim foveolatis, apice oblique recte truncatis, angulo suturali acuto (haud dehiscenti), exteriori dentato. ♂ ventris segmentis 3^o–5^m medio plagis duabus dense pilifero-punctatis, tibiisque quatuor posterioribus intus versus apicem pilosis, pectore nudo.

Long. 14–17 millim. ♂ ♀.

Hab. PANAMA, Volcan de Chiriqui (*Champion*).

The elytral foveoles are more uniform in size than in allied species, and all more or less round—with the exception of the usual impressed sutural stria, in which the ordinary small punctures are seen; the interstices are unequally spaced and nearly plane.

4. *Agra macracantha*.

Ab *A. castaneipedi* (♀) differt elytris apice suturali longe spinoso, foveolis plerumque minoribus, seriebus 2^a (a sutura), 6^a et 7^a versus apicem in striis impressis, interstitiis convexis.

Long. 17 millim. ♀.

Hab. PANAMA, Bugaba (*Champion*).

Possibly only a variety of *A. castaneipes*. In that species the sutural apex varies, being in some examples rectangular and in others slightly produced and acute. In *A. macracantha* it is lengthened into an acute spine, longer than that of the lateral angle of the truncature.

5. *Agra æneola*. (Tab. XII. fig. 13.)

A. mœrenti affinis et quoad formam similis. Gracilis, piceo-fusca, supra ænea vel cuprascens; antennis, partibus oris, pedibus piceo-rufis; capite sicut in *A. mœrenti* postice elongato, oblongo, lævi; thorace medio vix dilatato, antice gradatim angustato; elytris angustis, post medium paullo ampliatis, apice subrecte truncatis, sutura paullulum dehiscenti, angulo suturali nullomodo producto, exteriori breviter dentato, dorso lineatim foveolatis, foveolis subrotundatis, sat parvis, raro confluentibus.