

interstitio tertio tripunctato. Saturate olivaceo-ænea, antennis pedibusque chalybeo-nigris; elytrorum truncatura vix flexuosa, angulo suturali breviter, exteriori latius et longius, dentato; antennis articulo octavo haud abbreviato.

Long. 13 millim. ♀.

Hab. GUATEMALA, Teleman (*Champion*).

One example only.

19. *Agra mexicana*.

Agra mexicana, Buquet, Ann. Soc. Ent. Fr. 1835, p. 606¹; Chaudoir, ibid. 1861, p. 132.

Hab. MEXICO¹, Cordova (*Sallé*); GUATEMALA, Mirandilla, Teleman, Panzos (*Champion*).

Varies in its metallic coloration like *A. varians*, to which it is extremely closely allied. One example, from Panzos, is dark olivaceous green. The male has the whole centre of the mesosternum and abdomen, the inner side of the trochanters and femora, and the four posterior coxæ densely hairy, more hairy than in *A. varians* ♂, and resembling *A. cytherea* ♂.

20. *Agra chrysopteryx*. (Tab. XII. fig. 22.)

Agra chrysopteryx, Bates, Proc. Zool. Soc. 1878, p. 609¹.

"*A. rutilipenni* proxime affinis et similis. Castaneo-rufa, elytris (epipleuris exceptis) læte viridi-auratis; capite post oculos breviter semiovato, punctis nonnullis piliferis impresso; thorace medio modice rotundato-dilatato, paulo ante apicem fere subito angustato, lineatim subsparsim irregulariter foveato-punctato, longe piloso; elytris apice subrecte oblique truncatis, angulo exteriori breviter dentato, suturali rotundato, supra acute punctulato-striatis, striis secunda et quarta 5-6-punctatis, stria sexta bipunctata.

"♂ metasterno et abdomine medio longe rufo-barbatis.

"Long. 7 lin."

Hab. NICARAGUA, Chontales (*Janson*¹); PANAMA, Volcan de Chiriqui (*Champion*).

21. *Agra purpurea*. (Tab. XII. fig. 21.)

A. rutilipenni affinis. Cupreo-fusca, elytris (interdum olivaceo-æneis) marginibus læte igneo-cupreis, corpore subtus nigro-æneo nitido, antennis basi excepta piceo-rufis; capite post oculos modice elongato, ♀ quadrato, ♂ semiovato, occipite punctis nonnullis magnis; thorace sat lato, antice gradatim modice angustato, grosse lineato-punctato; elytris apice valde flexuoso-truncatis, sutura dehiscienti, angulo haud producto, angulo exteriori breviter dentato, punctulato-striatis, striis secunda, quarta et sexta punctis magnis et conspicuis plurimis.

♂ metasterno ventreque medio, trochanteribus et coxis 4 posterioribus femoribusque intermediis intus dense hirsutis.

♀ antennis articulo octavo haud abbreviato.

Long. 17 millim.

Hab. PANAMA, Volcan de Chiriqui (*Champion*).

The underside of the body and the epipleuræ of the elytra are shining, generally dark greenish brassy; the upper side is very dark purple-brown, the thorax more brightly metallic and the elytra sometimes greenish; the cincture of brilliant purplish copper which surrounds the elytra is limited on the sides to the marginal interstice,