

the basal and apical border being a little broader, and the suture being only very narrowly and sometimes indistinctly edged with the same colour.

22. *Agra ictina*.

A. purpureæ affinis, sed multo minor et gracilior, supra cupreo-fusca, versus elytrorum margines lætius rubro-cuprea, epipleuris viridi-æneis, antennis pedibusque piceis; capite post oculos modice elongato, ovato, occipite punctis magnis paucis; thorace medio sat rotundato, prope apicem angustato, dorso grosse punctato, punctis numerosis plerumque confluentibus, interspatiis lævibus, angustis, longitudinalibus; elytris angustis, apice flexuoso-truncatis, sutura dehiscenti angulo haud producto, angulo exteriori breviter dentato, punctato-striatis, striis secunda et quarta punctis majoribus sat conspicuis.

♂ metasterno ventrequè medio, coxis trochanteribusque 4 posterioribus intus, dense pilosis; femoribus fere nudis.

♀ antennis articulo octavo haud abbreviato.

Long. 12-14 millim.

Hab. MEXICO, Cordova (*Höge*); BRITISH HONDURAS, Rio Hondo (*Blancaneaux*); GUATEMALA, Teleman, Cubilguitz (*Champion*).

The colour of the elytra is dullish purple coppery, a little brighter sometimes on the second or third marginal interstice; but the margin of the elytra itself and the epipleura are always green. The thorax is sometimes bright brassy, and the punctuation, as in other species, is variable; it is, however, more crowded than usual in the allied species, the punctured stripe near the dorsal line consisting of punctures often three abreast, but very irregular.

23. *Agra auronitens*.

A. mexicana primo intuitu similis sed elytris læte viridi-auratis, antennis, palpis maxillaribus pedibusque rufis, geniculis fuscis; capite post oculos valde elongato quadrangulato, lævi; thorace sicut in *A. mexicana* punctis lineatis paucioribus; elytris apice utrinque valde bispinosis, punctulato-striatis, striis secunda et quarta punctis plurimis modice conspicuis.

♀ segmento ultimo ventrali medio apice profunde et anguste inciso; antennis articulo octavo præcedenti multo breviori.

Long. 15½ millim.

Hab. GUATEMALA, Teleman (*Champion*).

One example only.

24. *Agra panamensis*.

Agra panamensis, Bates, Proc. Zool. Soc. 1878, p. 609¹.

“Ad sectionem *A. æneæ* (F.) pertinet; at cæteris speciebus distinctissima. Latior, nigra, partibus oris, antennis, trochanteribus, genibus, tibiis et tarsis fulvo-castaneis; capite lato, pone oculos quadrato, margine posteriore ante collum exacte recto, supra impunctato; thorace quam caput longiore, pone medium rotundato-dilatato, antice usque ad apicem gradatim angustato, supra medio fere lævi, juxta marginem sparse punctato-impresso; elytris apice arcuatim truncatis, angulo exteriori fortiter spinoso, suturali acute dentato, supra vix inæqualibus, punctato-striatis, striis secunda et quarta seriatim rotundato-foveolatis, sexta et octava magis confuse foveolatis.

“♂ antennis articulo octavo multo abbreviato.

“Long. 12 lin.”

Hab. PANAMA (*Mr. E. Bartlett*).