

## PSEUDOMORPHA.

*Pseudomorpha*, Kirby, Trans. Linn. Soc. xiv. p. 98 (1825).

*Heteromorpha*, Kirby, *ibid.* t. 3. fig. 3.

*Drepanus*, Illiger, Mag. f. Insektenkunde, vi. p. 344 (1807), sine char.; Dejean, Sp. Gén. Col. v. p. 434 (1831).

*Axinophorus*, Dejean, Iconogr. i. p. 176.

Species of this genus have been recorded from Oceania and Australia as well as from America. It is doubtful, however, whether the Australian species really belong to the genus. Eight species have been described from widely different parts of America, North and South.

1. ***Pseudomorpha pilatei*.** (Tab. XII. fig. 25.)

*Pseudomorpha Pilatei*, Chaudoir, Rev. et Mag. Zool. 1862, p. 490<sup>1</sup>.

*Hab.* MEXICO, Yucatan (*Pilate*<sup>1</sup>).

A specimen of this species in the Sallé collection has a similar transverse pilose band in the middle of the fourth and fifth ventral segments which Dr. Horn mentions as distinguishing his *P. cronkhitei* of California.

2. ***Pseudomorpha behrensi*.**

*Pseudomorpha Behrensi*, Horn, Trans. Am. Ent. Soc. 1870, p. 76<sup>1</sup>.

*Hab.* NORTH AMERICA, California<sup>1</sup>.—MEXICO, Ciudad in Durango (*Forrer*).