

SUPPLEMENT.

DURING the progress of the foregoing work collections have been continually arriving containing new species, belonging, in many cases, to genera which had already been dealt with. These, together with a few others contained in the earlier collections, decision upon which was deferred until the arrival of further material, are described in the following Supplement, which will be final. Some synonymical notes and all the more important new localities of previously described species are added.

***Tetracha carolina*, var. *cyanides* (p. 1).**

I am informed by Mr. Flohr that the locality for this variety is the neighbourhood of Oaxaca, not Jalapa as stated. Herr Höge, in labelling his large collection on his return to Europe, attached by mistake the locality "Jalapa" to many of the Oaxaca species, and *vice versa* "Oaxaca" to species found at Jalapa.

CICINDELA (p. 4).

4 (A). ***Cicindela obsoleta*.**

Cicindela obsoleta, Say, Journ. Acad. Nat. Sci. Phil. iii. (1823), p. 143.

Var. Elytris subviolaceo-tinctis, immaculatis corporeque subtus cyaneo.

Hab. MEXICO, Ciudad in Durango (*Forrer*).

Agrees nearly with the typical form of the species from the Rocky Mountains in the colour of the upper surface; but the under surface is richly metallic blue, with here and there a greenish tinge, nearly as in the var. *vulturina*, Leconte.

7 (A). ***Cicindela euthales*.**

Cicindela euthales, Bates, Ann. & Mag. Nat. Hist. ser. 5, vol. ix. p. 319.

"*C. catharinæ* quoad formam similis, sed corporis lateribus nudis. Viridis, opaca; fronte, sutura lateribusque elytrorum nitidis; labro albo margine antico medio late producto tridentato, utrinque sinuato, angulis rectis; fronte verticali utrinque (et vertice) subtilissime striata; capite inter oculos haud concavo; palpis nigris; thorace transverso, lateribus albo pilosis vix rotundatis, antice angulatis, dorso transversim indistincte striguloso; elytris apice conjunctim rotundatis, dorso haud conspicue sculpturatis, immaculatis; corpore subtus viridi-nitido, lateribus cupreo-violaceis nudis; pedibus cupreis.

"*Var.* Nigra, opaca, abdomine medio et apice nitido; labro albo.

"Long. $4\frac{1}{2}$ – $5\frac{1}{2}$ lin. ♂ ♀.

"*Hab.* MEXICO, Ciudad in Durango (*Forrer*).

"Of similar form to *C. catharinæ*, but more closely allied to *C. ioessa* in the elytra not being visibly sculptured and in the naked sides of the body beneath. It differs from