

10 (c). **Cicindela lemniscata.**

Cicindela lemniscata, Leconte, Proc. Acad. Phil. 1854, p. 220¹.

Hab. NORTH AMERICA, Arizona, California¹.—MEXICO, Northern Sonora (*Morrison*).

20 (A). **Cicindela favergeri.**

Cicindela Favergeri, Audouin et Brullé, Archives du Muséum, i. p. 130, t. viii. fig. 6¹; Reiche, Revue Zool. 1842, p. 241².

Hab. PANAMA, San Feliz (*Champion*), Matashin (*Schunke*).—SOUTH AMERICA, Colombia^{1 2}, Ecuador.

Mr. Champion's examples belong to the variety in which the marginal white streak of the elytra is absent. Both forms are mentioned by Reiche as found together in Colombia, and Mr. Buckley took them both equally near Canelos, Ecuador.

25 (A). **Cicindela dugesi.**

C. punctulata similis, sed minor, lætius cuprea, abdomine rufo etc. Sat angusta, fuscescenti-rubro-cuprea, subnitida, subtus cum femoribus aurato-ænea, abdomine trochanteribusque omnibus rufis, tibiis rubro-cupreis, tarsis viridibus vel cupreis; elytris utrinque apud discum fasciola parum flexuosa obliqua guttaque paullo posteriori, albis; labro medio triangulariter producto (in ♂ minus quam in ♀) margine anteriori flexuoso dente parvo mediano.

Long. $3\frac{1}{2}$ – $4\frac{1}{4}$. ♂ ♀.

Hab. MEXICO, Tupataro, Morelia, Taretan (*Flohr*).

A distinct species of the *rufiventris* group recently sent me by Mr. Julius Flohr. In general form it is very similar to the common North-American *C. punctulata*, but it is of a redder coppery hue, and differs greatly in other ways. In shape of head and degree of prominence of eyes it resembles *C. rufiventris*, to which it is no doubt allied. The forehead is strongly strigose on each side, the occiput convex and irregularly sculptured. The thorax is more rounded in the middle of its sides than in *C. rufiventris* and allies, but its grooves are deep and blue-coloured as in those species, and the disk faintly transverse strigose. The elytra are narrower and more convex, but are similarly shaped at the apex, and have a minute sutural tooth; their surface is finely alutaceous, with minute shallow punctures set in greenish spots, and a row of larger ocellated punctures irregularly parallel to the suture, besides a shorter row of similar impressions in the basal depression on each side.

27 (A). **Cicindela guttifera.**

Cicindela guttifera, Leconte, Trans. Am. Phil. Soc. xi. 1856, p. 42¹.

Hab. NORTH AMERICA, New Mexico¹.—MEXICO, Northern Sonora (*Morrison*).

Generally considered to be a southern variety of the common North-American *C. duodecimguttata* (Dejean).

Cicindela lugens (p. 5).

Mr. Flohr informs me that Höge's specimens were not taken at Jalapa, but at