

Cicindela flohri (p. 9).

Jalapa as a locality again is wrong. The species does not occur, as Mr. Flohr informs me, out of the valley of Mexico.

Cicindela rectilatera (p. 10).

To the Mexican localities given add :—San Pedro Coahuila (*Dr. Palmer*).

Cicindela papillosa (p. 13).

To the localities given add :—PANAMA, Volcan de Chiriqui (*Champion*).

Some examples from this locality have a silky metallic green or golden lustre on the sides of the upper surface. Chontales examples are often slightly metallic ; in Mexico the insect is almost always sooty black above.

Cicindela viridisticta (p. 14).

Var. *arizonensis*. A typo differt elytrorum lateribus sicut dorso punctatis subnitidis, nec impunctatis, politis.

Hab. NORTH AMERICA, Arizona (*Morrison*).—MEXICO, Northern Sonora (*Morrison*).

Cicindela unicolor (p. 15).

Dr. Eugène Dugès informs Mr. Flohr that this species occurs in the Barranca de Beltran, on the road between Guadalajara and Colima, also near Guanajuato.

36 (A). **Cicindela castissima**. (Tab. XIII. fig. 1.)

C. signata (Dej.) affinis ; minor et angustior, subtus cum pedibus viridi-aurata, elytris albis sutura angusta (versus scutellum paullo dilatata) punctisque nonnullis medio basi viridi-æneis ; capite thoraceque cupreis, hoc creberrime sculpturato opaco, illo versus oculos nitidiori ; labro ♀ latissimo, brevi, antice fere recto edentato, albo ; thorace trapezoideo, lateribus quam in *C. signata* ♀ magis rectis ; elytris versus apicem oblique angustatis, apice ipso intus obliquato fere truncato ; thorace margine corporeque subtus lateribus dense albo-incumbenti-pilosis.

Long. $5\frac{1}{2}$ lin. ♀.

Hab. Arcas Islets, Gulf of Mexico.

45 (A). **Cicindela punctulata**.

Cicindela punctulata, Oliv. Ent. ii. 33, p. 27, t. 3. fig. 37.

Hab. MEXICO, Northern Sonora (*Morrison*).

Mr. Morrison sends one example of the green variety of this common North-American species.

ODONTOCHEILA (p. 15).

5 (A). **Odontocheila exilis**. (Tab. XIII. fig. 4.)

O. cinctulae simillima, forsan ejus varietas geographica. Paullo minor et gracilior ; elytrorum lateribus latius auratis, corporeque subtus toto cyaneo subviolaceo.

Long. $4\frac{1}{2}$ –5 lin. ♂ ♀.