

SCARITES (p. 30).

2. **Scarites subterraneus.**

Scarites subterraneus, Fabr. Mant. Ins. i. p. 206.

Var. Elytris fere parallelis, striis parum impressis, interstitiis subconvexis.

Hab. MEXICO, Northern Sonora (*Morrison*).

I place this var. under *Sc. subterraneus* rather than under *Sc. quadriceps*, owing to its more convex and posteriorly more abruptly declivous elytra. It approaches very near the *Sc. quadriceps*, var. *lissopterus*, from Texas.

DYSCHIRIUS (p. 31).

3. **Dyschirius analis.**

Dyschirius analis, Leconte, Ann. Lyceum N. York, v. p. 196¹.

Hab. NORTH AMERICA, River Colorado¹.—MEXICO, Northern Sonora (*Morrison*).

OXYDREPANUS (p. 31).

Oxydrepanus mexicanus (p. 31).

To the localities given add:—PANAMA, Tolé (*Champion*).

SCHIZOGENIUS (p. 37).

Schizogenius tristriatus (p. 38).

To the localities given add:—PANAMA, Volcan de Chiriqui (*Champion*).

Differs from Mexican typical individuals by the central pair of frontal carinae being thickened, and the three lateral ones on each side diminished in elevation and sharpness.

Schizogenius tenuis (p. 39).

To the localities given add:—PANAMA, Tolé (*Champion*).

7. **Schizogenius depressus.**

Schizogenius depressus, Leconte, Ann. Lyceum N. York, v. p. 197¹.

Hab. NORTH AMERICA, River Colorado¹.—MEXICO, Northern Sonora (*Morrison*).

PANAGÆUS (p. 41).

Panagæus sallæi (p. 41).

To the Mexican localities given add:—Chihuahua, Pinos Altos (*Hepburn*).

BIOL. CENTR.-AMER., Coleopt., Vol. I. Pt. 1, *July* 1884.

2 mm