

Chlænius ruficauda (p. 43).

To the Mexican localities given add:—Northern Sonora (*Morrison*).

Chlænius chlorochrous (p. 43).

To the localities given add:—PANAMA, San Feliz (*Champion*).

26. Chlænius regularis.

Chlænius regularis, Leconte, Ann. Lyceum N. York, v. p. 179¹.

Hab. NORTH AMERICA, California¹.—MEXICO, Northern Sonora (*Morrison*).

27. Chlænius glaucus.

Chlænius glaucus, Leconte, Proc. Ac. Phil. 1856, p. 28¹; Chaudoir, Monogr. des Chlæn. 1876, p. 239.

Hab. NORTH AMERICA, Junction of Gila and Colorado.—MEXICO, Northern Sonora (*Morrison*).

Chlænius tricolor (p. 44).

To the localities given add:—Northern Sonora (*Morrison*).

The Sonora form is similar to that found at Mexico city and Puebla, in which the head and thorax are coppery green.

Chlænius purpureus (p. 45).

To the Mexican localities given add:—Chihuahua, Mescales (*Hepburn*).

OODES (p. 45).

Oodes mexicanus (p. 45).

In the second part of a posthumous revised monograph of the *Oodides* (Ann. Soc. Ent. Fr. 1882, p. 496 *) Baron Chaudoir places this species in the genus *Stenocrepis*, to which his former genus *Crossocrepis* (including *O. mexicanus*) is subordinated as a section. Mr. Flohr has sent me a ♂ and ♀ of the species from Mexico, precise locality not stated, only 11 millim. long, and a third example (♂) 12½ millim., the same length as the ♀ example in the Sallé collection and the specimens described by Chaudoir.

Hab. NORTH AMERICA, Texas (*Chaudoir*).—MEXICO (*Sallé, Flohr*), Las Vigas (*Flohr*).

* The second part was published May 30th, 1883.