

STENOUS (p. 46).

Stenous duodecimstriatus (p. 46).

Add to the synonymy:—

Stenocrepis duodecimstriatus, Chaudoir, Monogr. des Oodides, Ann. Soc. Ent. Fr. 1882, p. 494¹.

Stenous Lecontei, Chaudoir, Bull. Mosc. 1857, ii. p. 41.

Oodes quatuordecimstriatus, Leconte, Ann. Lyc. N. York, iv. p. 331.

Oodes striatellus, Laferté, Ann. Soc. Ent. Fr. 1851, p. 272 (nota).

To the localities given add:—NORTH AMERICA, Texas, Louisiana¹.

This species belongs to the typical section of *Stenocrepis*, of which *Stenous* is a division in the revised monograph of Chaudoir above cited.

Stenous tibialis (p. 46).

Add to the synonymy:—

Oodes pallipes, Brullé, Voyage de D'Orbigny, vi. 2, t. 2. fig. 6¹.

To the localities given add:—SOUTH AMERICA, Bolivia¹.

S. chlorophanus, Erichson, is considered by Chaudoir to be distinct, though very closely allied.

Stenous gratiosus (p. 46).

Add as synonymy:—

Stenocrepis (Stenous) gratiosa, Chaudoir, Monogr. des Oodides, Ann. Soc. Ent. Fr. 1882, p. 500.

Stenous olivaceus (p. 47).

Chaudoir, in his revised monograph, is doubtfully inclined to consider this species as a variety of the South-American *S. metallicus*. The two are very closely allied, but appear to be as distinct as most others in this group.

STENOCREPIS (p. 47).

Stenocrepis chalcas (p. 47).

Stenocrepis chalcocrous, Chaudoir, Monogr. des Oodides, Ann. Soc. Ent. Fr. 1882, p. 487 (May 30, 1883).

POLYCHÆTUS (p. 47 ; to follow the genus *Stenocrepis*).

Polychætus, Chaudoir, Ann. Soc. Ent. Fr. 1882, p. 338.

Two species of this new genus are described by M. de Chaudoir, one Mexican and the other from Cayenne.

1. **Polychætus dejeani**.

Polychætus Dejeani, Chaudoir, l. c. p. 339.

Hab. MEXICO.