

Panama I have seen three males, whose tarsi agree in structure with those of Brazilian examples.

2. *Megadytes flohri*.

Megadytes flohri, Sharp, Trans. Roy. Dubl. Soc. n. s. ii. p. 709.

Hab. MEXICO, city of Mexico (*Flohr*), Vera Cruz (*Höge*), Cordova (*Sallé*).

The females in the small series of this species now before me are all destitute of any sexual sculpture, except that in two of them there exist ten or a dozen minute linear punctures clustered round the basal puncture of the internal series on the wing-case.

3. *Megadytes fraternus*.

Megadytes fraternus, Sharp, Trans. Roy. Dubl. Soc. n. s. ii. p. 708.

Hab. GUATEMALA, Dueñas, San Gerónimo (*Champion*).—ANTILLES, Guadeloupe, St. Domingo; SOUTH AMERICA, Demerara.

4. *Megadytes fallax*. (Tab. I. fig. 13.)

Cybister fallax, Aubé, Spec. Gén. vi. p. 54¹; Sharp, Trans. Roy. Dubl. Soc. n. s. ii. p. 710.

Hab. MEXICO (*Sallé*).—CAYENNE¹.

This little-known species is represented in Sallé's collection by a single female labelled "*Trogus flavocinctus*;" this determination, however, was erroneous, and the specimen, a very fine one, agrees with the two other females, these being all that we know of the species.

5. *Megadytes obesus*.

Megadytes obesus, Sharp, Trans. Roy. Dubl. Soc. n. s. ii. p. 710¹.

Hab. PANAMA¹.

6. *Megadytes giganteus*.

Cybister giganteus, Lap. Et. Ent. p. 99; Aubé, Spec. Gén. vi. p. 46; Sharp, Trans. Roy. Dubl. Soc. n. s. ii. p. 713¹.

Cybister lherminieri, Aubé, Spec. Gén. vi. p. 47.

Hab. ? MEXICO¹, Merida in Yucatan (*Sallé*); GUATEMALA, San Gerónimo (*Champion*); SALVADOR (*Sallé*).—BRAZIL; ANTILLES, Guadeloupe, Antigua, Cuba.

CYBISTER.

Cybister, Curt. Brit. Ent. iv. p. 151; Sharp, Trans. Roy. Dubl. Soc. n. s. ii. p. 918.

Trogus, Leach, Zool. Misc. iii. p. 70 (1817).

This important genus is especially characteristic of the eastern hemisphere, where it