

*Var. minor*, magis nitidus, paullo subtilius punctatus.

*Lispinus insularis*, Chevr. & Fauv. Ann. Soc. Ent. Fr. 1863, p. 442<sup>3</sup>; Notices ent. ii. p. 49<sup>4</sup>.

*Hab.* MEXICO (*Truqui*<sup>3 4</sup>); GUATEMALA, Zapote (*Champion*).—ANTILLES, Santo Domingo<sup>3 4</sup>, Cuba<sup>3 4</sup>.

I think it more than probable that *L. insularis* is only a small form of *L. granadensis*, though I have only seen one exponent, and that a not very satisfactory one, from our region; the larger form appears to be the least rare of our *Lispini*.

### 7. *Lispinus planus*.

*Lispinus linearis*, Fauv. (nec Er.), Bull. Soc. Linn. Normand. ix. p. 51<sup>1</sup>; Notices ent. ii. p. 47<sup>2</sup>.

*Lispinus planus*, Sharp, Trans. Ent. Soc. Lond. 1876, p. 416<sup>3</sup>.

*Hab.* MEXICO, Teapa (*Pilate*<sup>1 2</sup>).—SOUTH AMERICA, Colombia<sup>1 2</sup>, Amazons valley<sup>3</sup>.

A single example—one of Fauvel's types—is all I have seen from our region, and it is apparently conspecific with Amazonian specimens. In 'Notices ent.' vii. p. 13, Fauvel refers Leconte's *L. obscurus* to this species as a synonym, but from the remarks he makes it is probable it may be distinct.

### 8. *Lispinus anguinus*.

*Lispinus anguinus*, Fauv. Bull. Soc. Linn. Normand. ix. p. 54<sup>1</sup>; Notices ent. ii. p. 50<sup>2</sup>.

*Hab.* MEXICO, Teapa (*Pilate*<sup>1 2</sup>).—ANTILLES, Santo Domingo (*Sallé*<sup>1 2</sup>).

I have seen no example from our region of this distinct species.

### 9. *Lispinus fauveli*.

*Lispinus tenellus*, Chevr. & Fauv. (nec Er.), Ann. Soc. Ent. Fr. 1863, p. 443<sup>1</sup>; Fauv. Notices ent. ii. p. 53<sup>2</sup>.

Depressus, subnitidus, nigricans; prothorace, elytris antennisque piceis, harum basi pedibusque testaceis; abdominis apice rufescente; parcius omnino subtilissime punctulatus.

Long. 3 millim.

*Hab.* MEXICO<sup>1 2</sup>, Teapa (*Pilate*, in coll. Chevrolat), Cordova (*Sallé*); BRITISH HONDURAS, R. Hondo (*Blancaneaux*); GUATEMALA, near the city, Zapote, El Tumbador (*Champion*).—COLOMBIA<sup>1 2</sup>.—ANTILLES, Cuba<sup>1 2</sup>.

I do not think it at all probable that this insect is the species described by Erichson under the name of *L. tenellus*, and I therefore give it a new name. We have received altogether nearly two dozen examples of this insect, representing four races or possibly distinct species.