

This species is rather larger than *L. flavipennis* and has the thorax distinctly punctate and with a slender smooth space along the middle; it is equally close to *L. fauveli*, but is narrower, and has the thorax and elytra red, the former more distinctly punctate, the latter rather shorter. Eight examples.

#### 14. *Lispinus brevicollis*.

*Lispinus brevicollis*, Fauv. Bull. Soc. Linn. Normand. ix. p. 56<sup>1</sup>; Notices ent. ii. p. 52<sup>2</sup>.

*Hab.* MEXICO, Teapa<sup>1 2</sup>, Cordova (*Sallé*); GUATEMALA, near the city, El Jicaro, Cubilguitz, Cahabon (*Champion*); NICARAGUA, Granada (*Sallé*).

Of this rare species we have only single examples from each locality.

#### 15. *Lispinus longicornis*. (Tab. XIX. fig. 8.)

Rufus, depressus, pernitidus; antennis quam corpus vix brevioribus; parce punctatus; prothorace breve, basi utrinque longitudinaliter minus profunde impresso.

Long. 4 millim.

*Hab.* PANAMA, Volcan de Chiriqui 4000 to 6000 feet (*Champion*).

This insect is similar in form to *L. brevicollis*, but is distinguished not only by the very elongate antennæ but also by the very minute prosternal process separating the front coxæ. The head is broad and short, not margined in front; with prominent eyes, and very distinct widely distant antennal tubercles; sparingly punctate, and without foveæ. Thorax strongly transverse, a little narrower at the base, coarsely and rather irregularly punctate; not canaliculate, but with a narrow smooth space along the middle. Elytra very sparingly and obsoletely punctate, much longer than the thorax. Hind body also with a very minute almost invisible coriaceous sculpture. Unique.

#### 16. *Lispinus planellus*. (Tab. XIX. fig. 9.)

Depressus, rufo-testaceus, lævigatus, nitidissimus; prothorace ad basin constricto.

Long. 3 millim.

*Hab.* PANAMA, Volcan de Chiriqui 4000 to 6000 feet (*Champion*).

Antennæ short, moderately stout, piceous; joints 6-10 each shorter than long. Head short and broad, not margined, bi-impressed in front; antennary tubercles present, but flattened and broad. Thorax broader than long, flat; sides much rounded, narrowed and sinuate behind, and depressed close to the hind angles; with a very fine short channel along the middle, and the surface, though very polished, bearing a few very fine punctures. Elytra much longer than the thorax, polished and shining, with a single puncture on the disc of each.

This is an anomalous species, presenting to a great extent the appearance of an *Ancæus*; the front coxæ, however, are not at all prominent and the prosternum is developed into a very delicate process extending under and between them. We have received only one example.