

and deeply sculptured, with a broad groove along the middle in front. Elytra elongate, quite black, dull, with a conspicuous raised costa along the middle of each, sparingly and coarsely punctured. Hind body with the apical segments shining and obsoletely sculptured; basal segment dull, with some large depressions. Unique.

3. *Glyptoma cribellum*.

Piceum, opacum; capite nigricante undique fortiter punctato; abdomine æqualiter, densissime, fortissime profundeque punctato.

Long. $2\frac{1}{2}$ millim.

Hab. GUATEMALA, El Tumbador, Zapote, Capetillo (*Champion*); PANAMA, Volcan de Chiriqui (*Champion*).

Antennæ rather short, sparingly setose; joints 2–10 differing little from one another, strongly transverse. Head black, closely and deeply punctured, slightly shining. Thorax rugose, without depressions or channel. Elytra longer than the thorax, longitudinally rugose, with a well-marked costa along the middle of each. Hind body above and below, together with the breast, remarkably densely, deeply, and coarsely punctate-rugose; terminal segment very setose.

Of this species, so very distinct by the sculpture, Mr. Champion obtained a dozen examples at El Tumbador, and one each from the other localities; I regret to say that I lost the unique Panama exponent after determining it.

4. *Glyptoma hirticorne*. (Tab. XIX. fig. 11.)

Piceum, elytris rufescentibus, subnitidum; capite nigricante, polito, parce punctato; prothorace parvo, inæquali; elytris unicostatis; abdomine grosse punctato; antennis brevibus, medio dense pubescentibus. Long. $2\frac{1}{2}$ millim.

Hab. GUATEMALA, Capetillo (*Champion*).

Antennæ short; joints 2–10 strongly transverse, 3–6 densely, the others more sparingly, pubescent. Head with only a few distant fine punctures. Thorax narrower than the head, rounded in front, greatly narrowed behind, with two very large punctiform depressions in front of the base, and with some other large irregular punctures. Elytra elongate, red, rather shining, with a few coarse punctures, and along the middle of each an acutely raised costa. Hind body very coarsely sculptured, the sculpture consisting of irregular punctiform depressions which are coarser and deeper at the base of each segment. Five specimens.

5. *Glyptoma cognatum*.

Rufulum, sat nitidum; capite nigricante, polito, parce punctato; prothorace parvo, inæquali; elytris unicostatis; abdomine grosse punctato; antennis brevibus.

Long. $2\frac{1}{8}$ millim.

Hab. GUATEMALA, El Reposo (*Champion*).