

Hab. MEXICO, Vera Cruz (*Höge*).

Distinguished from *C. simplex* by the more oval form, but in other respects extremely similar to that species. A considerable series showing no variation has been received.

***Canthydrus rufipes* (p. 7).**

Before the localities given, insert:—MEXICO, Vera Cruz, San Juan, Bautista in Tabasco (*Höge*).

LACCOPHILUS (p. 9).

***Laccophilus apicalis* (p. 10).**

To the localities given, add:—PANAMA, Peña Blanca, Panama city (*Champion*).

***Laccophilus decipiens* (p. 10).**

To the Mexican locality given, add:—Chihuahua city (*Höge*).

In this, as well as in some other species, there occur rarely females with the epipleuræ dilated. I had previously supposed this to be a constant character for this sex in the species where I noticed it; but I am now convinced that in the case of this, and probably of other species, it is only a rare and exceptional occurrence: another very remarkable form of female dimorphism is thus added to those previously recognized as occurring in this family of Coleoptera.

***Laccophilus fuscipennis* (p. 10).**

The character of dilated epipleuræ is not a specific one, and this species, therefore, is distinguished from *L. confusus*, so far as I can at present judge, and we have received no more examples of it, solely by the larger size and the immaculate upper surface, and the absence of a coxal file in the female.

***Laccophilus confusus* (p. 11).**

To the Mexican locality given, add:—San Juan Bautista in Tabasco, Vera Cruz (*Höge*).

We have received now a good series of this species, and it appears at present to be really distinct from *L. americanus*: a form of the female with dilated epipleuræ occurs rarely. The coxal file is apparently always present in the female, though much more delicate than it is in the male.

10 (A). ***Laccophilus suffusus*.**

Laccophilus suffusus, Sharp, Trans. Roy. Dubl. Soc. n. s. ii. p. 293¹.

Hab. PANAMA, Bugaba (*Champion*).—SOUTH AMERICA, Bahia¹.