

may be known by the deeply emarginated and excavated labrum, and the form of the cephalic horn, which is elevated together with its closely-blended flanking tubercles.

### 1. *Verres corticicola*.

*Passalus corticicola*, Truqui, Rev. & Mag. Zool. 1857, p. 310<sup>1</sup>; Kaup, Harold's Col. Hefte, iv. p. 24.

*Verres corticicola*, Kaup, Monogr. p. 115, t. 7. f. 7<sup>2</sup>.

*Hab.* MEXICO<sup>1 2</sup>, Cordova (*Sallé*); GUATEMALA, Yzabal (*Sallé*), Cubilguitz, Coban, San Joaquin, Tactic (*Champion*); NICARAGUA, Chontales (*Janson*).

### 2. *Verres hageni*.

*Verres Hagenii*, Kaup, Monogr. der Passal. p. 116<sup>1</sup>.

*Hab.* MEXICO, Jalapa (*Höge*); GUATEMALA<sup>1</sup>, Capetillo (*Champion*); NICARAGUA, Chontales (*Janson*); COSTA RICA (*Sallé*, *Van Patten*).

### 3. *Verres cavicollis*. (Tab. I. figg. 20, 20 a.)

*V. furcilabro* affinis, labro apice profunde exciso et supra profunde et longe excavato; sed differt thoracis sulculo marginali antice utrinque in foveam latissimam haud profundam dilatato; clypeo antice valde flexuoso medioque profunde emarginato; cornu verticis cum tuberculis lateralibus consolidato, antice tricuspidato, dente mediano prolongato libero; carinis frontalibus a cornu separatis, arcum transversum parum curvatum efficientibus utrinque in tuberculum longe ante clypeum terminatum; metasterno punctulato-pubescenti, medio solum glabro, lateribus vitta punctulato-pubescenti, postice valde dilatata depressa, plagaque grossius punctata utrinque juxta apicem.

Long. 37 millim.; thorax 9, elytra 20 millim. long.

*Hab.* MEXICO, Jalapa (*Höge*).