

Toxpam, Jalapa, Matamoros Izucar, Parada, Orizaba, Guanajuato, Panistlahuaca (*Sallé*), Jalapa, Sayula, Misantla, Vera Cruz, Oaxaca, Rinconada, Cholula, Acapulco, Mazatlan, Tapachula (*Höge*), Mexico city (*Flohr*), Tehuantepec (*Sumichrast*); GUATEMALA (*Sallé*), San Gerónimo, Pantaleon, Mirandilla (*Champion*); NICARAGUA, Chontales (*Belt*).

Varies in size from 7 to 12 millim.; in colour from brilliant brassy-green and golden-coppery to bluish-green, violet, and deep black. In northerly localities the elytra are much less glossy. A small rotundate ovate form, with pygidium only slightly convex even in the male (*C. chloris*, *Sallé* coll.) occurs in various localities in Mexico.

### 25. *Canthon mutabilis*. (Tab. II. fig. 11, *var.*)

*Canthon mutabile*, Lucas, in Voy. de Castelnau, Zool. iii. p. 100<sup>1</sup>.

*Canthon mutabilis*, Harold, Berl. ent. Zeitschr. 1868, p. 127<sup>2</sup>.

*Hab.* PANAMA, Bugaba, Volcan de Chiriqui, Tolé (*Champion*).—SOUTH AMERICA<sup>2</sup>, Colombia<sup>2</sup>, Amazons<sup>1 2</sup>, La Plata<sup>2</sup>.

The Panama examples form a local variety of the species similar to the *var. transversalis* described by Von Harold, but differing in the bright brassy-green head and thorax. In Colombia the local form is similar, with the exception of having a brassy-black head and thorax. Some examples from the Lower Amazons are like the Colombian form, but they are there mingled with a great diversity of colour-variations.

An example from Tolé is figured.

### 26. *Canthon deyrollei*.

*Canthon deyrollei*, Harold, Berl. ent. Zeitschr. 1868, p. 132<sup>1</sup>.

*Hab.* BRITISH HONDURAS, R. Sarstoon (*Blancaneaux*); GUATEMALA, San Gerónimo (*Champion*); NICARAGUA, Chontales (*Belt*).

The locality of the species was unknown to Von Harold<sup>1</sup>. *C. deyrollei* strongly resembles the blue-black colour-variety of *C. chevrolati*, from which its pluridentate clypeus at once distinguishes it.

### 27. *Canthon chlorizans*.

Obscure viridi-metallicus, nitidus, pedibus rufo-testaceis; capite thoraceque sat crebre sed discrete punctatis; clypeo æqualiter quadri-dentato, sutura genarum haud dentata; elytris sat acute simpliciter striatis, interstitiis disperse punctulatis, humeris haud carinulatis; pygidio parum convexo, punctato, medio prope basin obtuse carinato; corpore subtus nigro, subopaco; tibiis posticis subrectis haud dilatatis.

♂. Tibiarum anticarum calcar apice lato bifido.

Long. 5 millim.

*Hab.* MEXICO, Juquila (*Höge*).

A little more oblong, and with the elytra relatively longer, than *C. viridis*.