

♂. Caput subplanum, disperse punctulatum; clypeo elongato, apice late reflexo et sinuato; vertice cornubus duobus semicirculariter curvatis, cupreis, basi haud carina connexis. Thorax dorso antice obtuse lobatus, lobo post apicem constricto. Tibiæ anticæ valde elongatæ, curvatæ; apice intus prolongato vix reflexo, penicillato.

Long. $8\frac{1}{2}$ millim. ♂.

Hab. PANAMA, San Feliz (*Champion*).

8. *Onthophagus rhinolophus*. (Tab. V. figg. 5, ♂; 5 a, side view of head and thorax.)

Onthophagus rhinolophus, Harold, Ann. Soc. Ent. Fr. 1869, p. 510¹.

Hab. MEXICO, San Andres Tuxtla (*Boucard*¹, in coll. *Sallé*), Oaxaca (*Höge*); GUATEMALA, San Isidro (*Champion*); PANAMA, Volcan de Chiriqui (*Champion*).

The single Guatemalan example, a female, differs in its rather more deeply-impressed and more strongly-punctured striæ. In the very finely, scarcely visibly, punctured elytral interstices it agrees with the typical specimens. We figure a male example from Tuxtla.

9. *Onthophagus belorhinus*. (Tab. V. figg. 6, ♂; 6 a, side view of head and thorax; 7, ♀; 7 a, side view of head and thorax.)

O. rhinolophus proxime affinis, a quo differt colore fusco-æneo nitido, thorace grosse punctato, elytris grosse crenato-punctatis; thorace (♂) antice medio lobato, lobo longitudinaliter sulcato; clypeo (♂) sicut in *O. rhinolophus*, late truncato et medio cornu erecto armato; genis margine explanato-reflexo, recto, juxta clypei suturam valde angulato, vertice bicornuto. ♀ clypeo triangulari sat elongato, apice bidentato, supra punctato-rugoso, antice transversim carinata, vertice breviter bicornuto; thorace antice medio tuberculis duobus compresso-conicis.

♂ major. Cornu clypei basi gracili, apice valde et abrupte dilatato, cornubusque verticis thorace altioribus.

♂ minor. Cornu clypei basi latior apice obtuso nec dilatatus; cornubus verticis sicut in ♂ majore.

♂ effœminatus. Clypeo apice leviter producto, reflexo, apice emarginato; vertice bituberculato thoraceque antice medio haud lobato, bituberculato.

♀ minor. Vertex bituberculatus, thorace tuberculis obsoletis.

Long. 5–7 $\frac{1}{2}$ millim.

Hab. GUATEMALA, El Reposo, Las Mercedes, San Isidro, Pantaleon, Zapote (*Champion*).

Var.? ♂ minor. Viridi-auratus, elytris cupreis, pedibus rufis.

Hab. NICARAGUA, Chontales (*Janson*).

Many examples of the typical form, one only of the variety.

This and the preceding species belong to a small group* of the genus in which both

* This curious and beautiful little group appears to be peculiar to Tropical America. Other species belonging to it are *O. clypeatus* (Blanch.), Bolivia, *O. rhinophyllus* (Har.), Venezuela and Colombia, and the two following:—

1. *Onthophagus xanthomerus*.

Niger, nitidus, elytris subviolaceis, sericeo-opacis, femoribus flavis; thorace punctulato; elytris subtiliter punctato-striatis, interstitiis planis, alutaceis, punctulatis. ♂ minor(?) clypeo subtriangulari, medio cornu erecto,