

14. **Onthophagus nasicornis.** (Tab. V. figg. 13, ♂ ; 13 a, side view of head and thorax.)

Onthophagus nasicornis (Dej. Cat.), Harold, Ann. Soc. Ent. Fr. 1869, p. 511¹.

Nigerrimus, politus ; thorace sparsim punctulato ; elytrorum striis punctato-crenatis, interstitiis parum convexis, sublævibus ; pygidio crebre punctato.

♂. Clypeus subtriangularis, medio apice cornu erecto laminiformi apice haud dilatato ; vertice haud cornuto. Thorax antice verticalis tri-impressus, dorso anteriore quadrato-gibboso.

♀. Clypeus bidentatus vel apice profunde fissus ; carina frontalis brevis, transversa. Thorax convexus.

Hab. MEXICO¹, Cordova, Tuxtla (*Sallé*).

The head in the male is narrower than in the preceding species, the genæ not being greatly dilated, and their outline continuous with that of the clypeus up to the base of the horn.

15. **Onthophagus dicranus.** (Tab. V. figg. 14, ♂ ; 14 a, side view of head and thorax.)

Parvus, fusco-niger, subnitidus, cupreo-tinctus, pedibus rufo-piceis ; capite thoraceque grosse sat confertim punctatis ; elytrorum striis valde crenato-punctatis, interstitiis rugulosis et sparsim punctatis.

♂ major. Clypeus apice cornu erecto laminato furcato armatus, apice obtuse truncatus, margine leviter flexuoso. Thorax dorso anteriore medio bituberculato, tuberculis altis, compresso-conicis.

♂ minor. Cornu clypei abbreviatum, apice obtusum.

♀. Clypeus bidentatus, capite supra plano, carina frontali brevissima, obsoleta. Long. 5-6 millim.

Hab. PANAMA, Bugaba (*Champion*).

The two conical tubercles of the thorax in fully developed males are greatly elevated and porrected, forming a bicuspid protuberance which overhangs the hind margin of the head. In the minor males they are scarcely more elevated than the thoracic tubercles of *O. rhinolophus*.

16. **Onthophagus mexicanus.** (Tab. V. figg. 15, ♂ ; 15 a, side view of head and thorax.)

O. latebroso (Fabr.) affinis, sed differt corpore supra fere calvo etc. Subopacus, atro-cæruleus vel violaceus, interdum viridis, raro cupreo-auratus ; thorace subtiliter disperse punctulato, medio basi læviore ; elytris subtilissime punctulato-striatis, interstitiis planissimis, punctulatis, punctulis brevissime setiferis ; antennis nigro-clavatis ; clypeo (♂ ♀) medio apice obtuso truncato, sinuato, supra ♂ lævi, ♀ punctato-rugoso.

♂ major. Caput margine utrinque valde flexuoso bilobato, clypeo lato et valde reflexo ; carinis transversis duabus, anteriore subtili transversa, posteriore arcuata utrinque in cornu acuto producta. Thorax antice verticalis profunde bi-impressus ; dorso medio antice valde prolongato, furcato. Tibiis anticis elongatis, apice intus acute productis.

♂ effoeminatus. Capitis margine laterali minus flexuoso, carinis validioribus, posteriore utrinque haud prolongata. Thorax dorso antice parum producto, quadrato-lobato, lobo antice leviter emarginato. Tibiis anticis apice intus acutis.

♀. Fere sicut in ♂ effoeminato ; differt thoracis lobo anteriore minus producto haud emarginato tibiisque anticis intus apice simplicibus.

Long. 7-10 millim.